

QUIVER VARIETIES AND CLUSTER ALGEBRAS

HIRAKU NAKAJIMA

ABSTRACT. Motivated by a recent conjecture by Hernandez and Leclerc [31], we embed a Fomin-Zelevinsky cluster algebra [21] into the Grothendieck ring $\mathbf{R}$ of the category of representations of quantum loop algebras $\mathbf{U}_q(\mathbf{Lg})$ of a symmetric Kac-Moody Lie algebra, studied earlier by the author via perverse sheaves on graded quiver varieties [49]. Graded quiver varieties controlling the image can be identified with varieties which Lusztig used to define the canonical base. The cluster monomials form a subset of the base given by the classes of simple modules in $\mathbf{R}$, or Lusztig's dual canonical base. The positivity and linearly independence (and probably many other) conjectures of cluster monomials [21] follow as consequences, when there is a seed with a bipartite quiver. Simple modules corresponding to cluster monomials factorize into tensor products of 'prime' simple ones according to the cluster expansion.

CONTENTS

1. Introduction	1
2. Preliminaries (I) – Cluster algebras	7
3. Preliminaries (II) – Graded quiver varieties	11
4. Graded quiver varieties for the monoidal subcategory $\mathcal{C}_1$	20
5. From Grothendieck rings to cluster algebras	24
6. Cluster character and prime factorizations of simple modules	26
7. Cluster algebra structure	34
References	40

1. INTRODUCTION

1.1. Cluster algebras. Cluster algebras were introduced by Fomin and Zelevinsky [21]. A cluster algebra $\mathcal{A}$ is a subalgebra of the rational function field $\mathbb{Q}(x_1, \dots, x_n)$ of n indeterminates equipped with a distinguished set of variables (cluster variables) grouped into overlapping subsets (clusters) consisting of n elements, defined by a recursive procedure (mutation) on quivers. Let us quote the motivation from the original text [loc. cit., p.498, the second paragraph]:

This structure should serve as an algebraic framework for the study of “dual canonical bases” in these coordinate rings and their q -deformations. In particular, we conjecture that all monomials in the variables of any given cluster (the cluster *monomials*) belong to this dual canonical basis.

Here “dual canonical base” means a conjectural analog of the dual of Lusztig canonical base of $\mathbf{U}_q^-$, the $-$ part of the quantized enveloping algebra ([43]). One of deepest properties of the dual canonical base is positivity: the structure constants are in $\mathbb{Z}_{\geq 0}[q, q^{-1}]$. But the existence and positivity are not known for cluster algebras except some examples.

2000 *Mathematics Subject Classification.* Primary 17B37; Secondary 14D21, 16G20.

Supported by the Grant-in-aid for Scientific Research (No.19340006), JSPS.

The theory of cluster algebras has been developed in various directions different from the original motivation. See the list of references in a recent survey [37].

One of most active directions is the theory of the cluster category [6]. It is defined as the orbit category of the derived category $\mathcal{D}(\text{rep } \mathcal{Q})$ of finite dimensional representations of the initial quiver $\mathcal{Q}$ under the action of an automorphism. This theory is quite useful to understand combinatorics of the cluster algebra: clusters are identified with tilting objects, and mutations are interpreted as exchange triangles. We refer to the survey [37] again.


However the cluster category *does not* have enough structures, compared with the cluster algebra. For example, multiplication of the cluster algebra roughly corresponds to the direct sum of the cluster category, but addition remains obscure. So the cluster category is called *additive categorification* of the cluster algebra. The cluster algebra is recovered from the cluster category by the so-called cluster character. (Somebody calls Caldero-Chapoton map.) But it is not clear how to obtain all the “dual canonical base” elements from this method.

Very recently Hernandez and Leclerc [31] propose another categorical approach. They conjecture that there exists a monoidal abelian category $\mathcal{M}$ whose Grothendieck ring is the cluster algebra. All of structures of the cluster algebra can be conjecturally lifted to the monoidal category. For example, the dual canonical base is given by simple objects, the combinatorics of mutation is explained by decomposing tensor products into simple objects, etc. Here we give the table of structures:

cluster algebra	additive categorification	monoidal categorification
$+$	$?$	$\oplus$
$\times$	$\oplus$	$\otimes$
clusters	cluster tilting objects	real simple objects
mutation	exchange triangle	$0 \rightarrow S \rightarrow X_i \otimes X_i^* \rightarrow S' \rightarrow 0$
cluster variables	rigid indecomposables	real prime simple objects
dual canonical base	$?$	simple objects
$?$	$?$	prime simple objects

In the bottom line, we have a definition of *prime simple* objects, those which cannot be factored into smaller simple objects. There is no counter part in the theory of the cluster algebra, so completely new notion.

However the monoidal categorification seems to have drawback. We do not have many tools to study the tensor product factorization in abstract setting. We need an additional input from other sources. Therefore it is natural to demand functors connecting two categorifications exchanging $\oplus$ and $\otimes$, and hopefully ‘?’ and $\oplus$. We call them *tropicalization* and de-tropicalization functors¹ expecting the top ? in the additive categorification column is something like ‘min’:



The author believes this is an interesting idea to pursue, but it is so far just a *slogan*: it seems difficult to make even definitions of (de)tropicalization functors precise. Therefore we set aside categorical approaches, return back to the origin of the cluster algebra, i.e. the construction of the canonical base, and ask why it has many structures ?

¹Leclerc himself already had a hope to make a connection between two categorifications ([40]). He calls ‘exponential’ and ‘log’.

The answer is simple: Lusztig's construction of the canonical base is based on the category of perverse sheaves on the space of representations $\mathbf{E}_W$ of the quiver. Therefore

- (a) it has the structure of the monoidal abelian category, where the tensor product is given by the convolution diagram coming from exact sequences of quiver representations;
- (b) it inherits various combinatorial structure from the module category $\text{rep } \mathcal{Q}$, and probably also from the cluster category.

In this sense, we already have (de)tropicalization functors !

Thus we are led to ask a naive question, sounding much more elementary compared with categorical approaches:

Is it possible to realize a cluster algebra entirely in Lusztig's framework, i.e. via a certain category of perverse sheaves on the space $\mathbf{E}_W$ of representations of a quiver ?

If the answer is affirmative, the positivity conjecture is a direct consequence of that of the canonical base.

As far as the author searches the literature in the subject, there is no explicit mention of this conjecture, though many examples of cluster algebras arise really as subalgebras of $\mathbf{U}_q^-$. Usually Lusztig's perverse sheaves appear only as a motivation, and is not used in a fundamental way. A closest result is Geiss, Leclerc and Schröer's work [25, 26] where the cluster algebra is realized as a space of constructible functions on Λ_W , the space of nilpotent representations of the preprojective algebra. This Λ_W is a lagrangian in the cotangent space $T^*\mathbf{E}_W$ of the space $\mathbf{E}_W$ of representations. The space of constructible functions was used also by Lusztig to construct the *semicanonical* base [45]. Constructible functions vaguely related to perverse sheaves (or D -modules) via characteristic cycle construction, though nobody makes the relation precise. And it was proved that cluster monomials are indeed elements of the dual semicanonical base [25, 26]. But constructible functions have less structures than perverse sheaves, in particular, the positivity of the multiplication is unknown.

Now we come to a point to explain the place where the author looks for the candidate of the framework. It is another geometric construction of an algebra together with distinguished (canonical) base in author's earlier work [49]. It is another child of Lusztig's work.

1.2. Graded quiver varieties and quantum loop algebras. In [49] the author studied the category $\mathcal{R}$ of l -integrable representations of the quantum loop algebra $\mathbf{U}_q(\mathbf{Lg})$ of a symmetric Kac-Moody Lie algebra $\mathbf{g}$ via perverse sheaves on graded quiver varieties $\mathfrak{M}_0^\bullet(W)$ (denoted by $\mathfrak{M}_0(\infty, \mathbf{w})^A$ in [loc. cit.]). If $\mathbf{g}$ is a simple Lie algebra of type ADE , $\mathbf{U}_q(\mathbf{Lg})$ is a subquotient of Drinfeld-Jimbo quantized enveloping algebra of affine type ADE (usually called the quantum affine algebra), and $\mathcal{R}$ is nothing but the category of finite dimensional representations of $\mathbf{U}_q(\mathbf{Lg})$. The graded quiver varieties are fixed point sets of the quiver varieties $\mathfrak{M}_0(W)$ introduced in [47, 48] with respect to torus actions. The main result says that the Grothendieck group $\mathbf{R}$ of $\mathcal{R}$ has a natural t -deformation $\mathbf{R}_t$ which can be constructed from a category $\mathcal{P}_W$ of perverse sheaves on $\mathfrak{M}_0^\bullet(W)$ so that simple (resp. standard) modules correspond to dual of intersection cohomology complexes (resp. constant sheaves) of natural strata of $\mathfrak{M}_0^\bullet(W)$. Here the parameter t comes from the cohomological grading. Furthermore the transition matrix of two bases of simple and standard modules (= dimensions of stalks of IC complexes) is given by analog of Kazhdan-Lusztig polynomials, which can be computed² by using purely combinatorial objects $\chi_{q,t}$, called t -analog of q -characters [51, 54]. If we set $t = 1$, we get the

²The meaning of the word *compute* will be explained in Remark 6.4.

q -character defined by [38, 24] as the generating function of the dimensions of l -weight spaces, simultaneous generalized eigenspaces with respect to a commutative subalgebra of $U_q(\mathbf{Lg})$. For the simple module corresponding to an IC complex L , $\chi_{q,t}$ is the generating function of multiplicities of L in direct images of constant sheaves on various nonsingular graded quiver varieties $\mathfrak{M}^\bullet(V, W)$ under morphisms $\pi: \mathfrak{M}^\bullet(V, W) \rightarrow \mathfrak{M}_0^\bullet(W)$.

We have a noncommutative multiplication on $\mathbf{R}_t$, which is a t -deformation of a commutative multiplication on $\mathbf{R}$. When $\mathbf{g}$ is of type ADE , the commutative multiplication on $\mathbf{R}$ comes from the tensor product $\otimes$ on the category $\mathcal{R}$ as $U_q(\mathbf{Lg})$ is a Hopf algebra. (It is not known whether the quantum loop algebra $U_q(\mathbf{Lg})$ can be equipped with the structure of a Hopf of algebra in general.) The t -deformed multiplication was originally given in terms of t -analog of q -characters, but Varagnolo-Vasserot [58] later introduced a convolution diagram on $\mathfrak{M}_0^\bullet(W)$ which gives the multiplication in more direct and geometric way.

These geometric structures are similar to ones used to define the canonical base of U_q^- by Lusztig [43]. We have the following table of analogy:

$\mathbf{R}_t$	geometry	dual of U_q^-
standard modules $M(W)$	constant sheaves	dual PBW base elements
simple modules $L(W)$	IC complexes	dual canonical base elements
t -deformed $\otimes$	convolution diagram	multiplication

Note that U_q^- is not commutative even at $q = 1$, while its dual $(U_q^-)^*|_{q=1}$ is the coordinate ring $\mathbb{C}[\mathfrak{n}^-]$, hence commutative. Hence we should compare $\mathbf{R}_t$ with $(U_q^-)^*$, not with U_q^- . Also the convolution diagram looks similar to one for the comultiplication, not to one for the multiplication. The only difference is relevant varieties: Lusztig used the vector spaces $\mathbf{E}_W$ of representations of the quiver with group actions (or the moduli *stacks* of representations of the quiver), while the author used graded quiver varieties, which are framed moduli *spaces* of graded representations of the preprojective algebra associated with the underlying graph.

The computation of the transition matrix is hard to use in practice, like the Kazhdan-Lusztig polynomials. On the other hand many peoples have been studying special modules (say tame modules, Kirillov-Reshetikhin modules, minimal affinization, etc.) by purely algebraic approaches, at least when $\mathbf{g}$ is of finite type. See [11] and the references therein. Their structure is different from that of general modules. Thus it is natural to look for a special geometric property which holds *only* for graded quiver varieties corresponding to these classes of modules. In [49, §10] the author introduced two candidates of such properties. We name corresponding modules *special* and *small* respectively. These properties are easy to state both in geometric and algebraic terms, but it is difficult to check whether a given module is special or small. Since [loc. cit.], we have been gradually understanding that smallness is not a right concept as there are only very few examples (see [30]), but the speciality is a useful concept and there are many special modules, say Kirillov-Reshetikhin modules³. One of applications of this study was a proof of the T -system, which was conjectured by Kuniba-Nakanishi-Suzuki in 1994 (see [53] and the references therein). Several steps in the proof of the main result in [loc. cit.] depended on the geometry, but they were replaced by purely algebraic arguments, and generalized to nonsymmetric quantum loop algebras cases later by Hernandez [29]. It was a fruitful interplay between geometric and algebraic approaches.

³Special modules form a *special* class of modules. Small modules form a *small* class of modules. But the name ‘small’ originally comes from the smallness of a morphism.

1.3. Realization of cluster algebras via perverse sheaves. Hernandez and Leclerc [31] not only give an abstract framework of the monoidal categorification of the cluster algebra, but also its candidate. It is a certain monoidal (i.e. closed under the tensor product) subcategory $\mathcal{C}_1$ of $\mathcal{R}$ when $\mathfrak{g}$ is of type ADE . They indeed show that $\mathcal{C}_1$ is a monoidal categorification for type A and D_4 . Therefore we have a strong evidence that it is a right candidate. From what we have reviewed just above, if it indeed is a monoidal categorification, the cluster algebra is a subalgebra of $\mathbf{R}$, constructed via perverse sheaves on graded quiver varieties ! Moreover, from the philosophy explained above, we could expect that graded quiver varieties corresponding to $\mathcal{C}_1$ have very *special* features compared with general ones.

In this paper, we show that it turns out to be true. The first main observation (see Proposition 4.6) is that the graded quiver varieties $\mathfrak{M}_0^*(W)$ become just the vector spaces $\mathbf{E}_W$ of representations of the *decorated quiver*. Here the decorated quiver⁴ is constructed from a given finite graph with a bipartite orientation by adding a new (frozen) vertex i' and an arrow $i' \rightarrow i$ (resp. $i \rightarrow i'$) if i is a sink (resp. source) for each vertex i . (See Definition 4.3.) Therefore the underlying variety is nothing but what Lusztig used. Also the convolution diagram turns out to be the same as Lusztig's one. Thus the Grothendieck group $\mathcal{K}(\mathcal{C}_1)$ is also a subalgebra of the dual of $\mathbf{U}_q^-$, associated with the Kac-Moody Lie algebra corresponding to the decorated quiver.

To define a cluster algebra with frozen variables (or with coefficients in [21]), we choose a quiver with choices of frozen vertexes. We warn the reader that this quiver for the cluster algebra (we call $\mathbf{x}$ -quiver, see Definition 5.4) is slightly different from the decorated quiver: the principal part has the opposite orientation while the frozen part is the same.

1.4. Second key observation. Once we get a correct candidate of the class of perverse sheaves, we next study structures of the dual canonical base and try to pull out the cluster algebra structure from it. We hope to see a shadow of the structure of a cluster category.

The dual of the subalgebra is a quotient. Thus we introduce an equivalence relation on the canonical base. The second key observation is that each equivalence class contains an exactly one skyscraper sheaf $1_{\{0\}}$ of the origin 0 of $\mathbf{E}_W$ (the simplest perverse sheaf !). This equivalence relation is built in the theory of graded quiver varieties. From this observation together with the first observation that the graded quiver varieties are vector spaces, we can apply the Fourier-Sato-Deligne transform [36, 39] to make a reduction to a study of constant sheaves $1_{\mathbf{E}_W^*}$ on the whole space.

There is a certain natural family of projective morphisms $\pi^\perp: \tilde{\mathcal{F}}(\nu, W)^\perp \rightarrow \mathbf{E}_W^*$ from non-singular varieties $\tilde{\mathcal{F}}(\nu, W)^\perp$. This family appears as monomials in Lusztig's context, and q -characters in the theory reviewed in §1.2. Using these morphisms, we define a homomorphism from $\mathbf{R}$ to the cluster algebra. Fibers of these morphisms are what are called *quiver Grassmannian* varieties. People study their Euler characters and define the cluster character as their generating function. This is clearly related to the study of the pushforward

$$\pi_!^\perp(1_{\tilde{\mathcal{F}}(\nu, W)^\perp}[\dim \tilde{\mathcal{F}}(\nu, W)^\perp]).$$

If $\mathbf{E}_W^*$ contains an open orbit, then the Euler number of the fiber over a point in the orbit is nothing but the coefficient of $1_{\mathbf{E}_W^*}[\dim \mathbf{E}_W]$ in the above push-forward. When the dual canonical base element is a cluster monomial, $\mathbf{E}_W^*$ indeed contains an open orbit. Therefore we immediately see that all cluster monomials are dual canonical base elements. This very

⁴The decorated quiver is different from one in [46], where there are no arrows between i and i' .

simple observation between the cluster character and the push-forward was appeared in the work of Caldero-Reineke [9]⁵.

To be more precise, we need to apply reflection functors at all sink vertexes in the decorated quiver with opposite orientations to identify fibers of $\tilde{F}(\nu, W)^\perp$ with quiver Grassmannian varieties. The resulted quiver corresponds to the cluster algebra with principal coefficients [23].

An appearance of the cluster character formula in the category $\mathcal{C}_1$ was already pointed out in [31, §12], as it is nothing but a leading part of the q -character mentioned above. (We call the leading part the *truncated* q -character.)

From a result on graded quiver varieties, it also follows that quiver Grassmannian varieties have vanishing odd cohomology groups under the above assumption. The generating function of all Betti numbers is nothing but the truncated t -analog of q -character of a simple modules. In particular, it was computed in [54].

But the only necessary assumption we need is that perverse sheaves corresponding to canonical base elements have strictly smaller supports than $\mathbf{E}_W^*$ except $1_{\mathbf{E}_W^*}[\dim \mathbf{E}_W^*]$. Even if this condition is not satisfied, we can consider the *almost simple module* $\mathbb{L}(W)$ corresponding to the sum of perverse sheaves whose supports are the whole $\mathbf{E}_W^*$. Then the total sum of Betti numbers (the Euler number is not natural in this wider context) of the quiver Grassmannian give the truncated q -character of the almost simple module. An almost simple module $\mathbb{L}(W)$ is not necessarily simple in general.

It is rather simple to study tensor product factorization of $\mathbb{L}(W)$ since we computed their truncated q -characters. First we observe that Kirillov-Reshetikhin modules simply factor out. Then we may assume that W have 0 entries on frozen vertexes. Thus W is supported on the first given vertexes. We next observe that $\mathbb{L}(W)$ factors as

$$\mathbb{L}(W) \cong \mathbb{L}(W^1) \otimes \cdots \otimes \mathbb{L}(W^s),$$

according to the *canonical decomposition* $W = W^1 \oplus \cdots \oplus W^s$ of W . Recall the canonical decomposition is the decomposition of a generic representation of $\mathbf{E}_W$ first introduced by Kac [34, 35], and studied further by Schofield [56]. It is known that each W^k is a Schur root (i.e. a general representation is indecomposable) and Ext^1 between generic representations from two different factors W^k, W^l vanish.

We prove that a simple module $L(W)$ corresponds to a cluster monomial if and only if the canonical decomposition contains only real Schur roots. In this case, we have $\mathbb{L}(W) = L(W)$, $\mathbb{L}(W^k) = L(W^k)$ and each $L(W^k)$ corresponds to a cluster variable, and the above tensor factorization corresponds to the cluster expansion.

1.5. To do list. In this paper, basically due to laziness of the author, at least four natural topics are not discussed:

- Our Grothendieck ring $\mathbf{R}$ has a natural noncommutative deformation $\mathbf{R}_t$. It should contain the quantum cluster algebra in [4]. In fact, we already give our main formula (in Theorem 6.3) in Poincaré polynomials of quiver Grassmannian varieties. Therefore the only remaining thing is to prove the quantum version of the cluster character formula. Any proof in the literature should be modified to the quantum version naturally, as it is based on counting of rational points.

⁵There is a gap in the proof of [9, Theorem 1] since Lusztig's v is identified with q . The correct identification is $v = -\sqrt{q}$.

- We only treat the case when the underlying quiver is bipartite. Since the choice of the quiver orientation is not essential in Lusztig's construction (in fact, the Fourier transform provides a technique to change orientations), this assumption probably can be removed.
- We only treat the symmetric cases. Symmetrizable cases can be studied by considering quiver automorphisms as in Lusztig's work. Though the corresponding theory was not studied in author's theory, it should corresponds to the representations of twisted quantum affine algebras.
- In [25, 26] it was proved that cluster monomials are semicanonical base elements. It was conjectured that they are also canonical base elements. It is desirable to study the precise relation of this work to ours.

The author or his friends will hopefully come back to these problems in near future.

In [31] a further conjecture is proposed for the monoidal subcategory $\mathcal{C}_\ell$, where $\mathcal{C}_1$ is the special case $\ell = 1$. Since the graded quiver varieties are no longer vector spaces for $\ell > 1$, the method of this paper does not work. But it is certainly interesting direction to pursue. We also remark that other connections between the cluster algebra theory and the representation theory of quantum affine algebras have been found by Di Francesco-Kedem [18] and Inoue-Iyama-Kuniba-Nakanishi-Suzuki [33]. It is also interesting to make a connection to their works.

This article is organized as follows. §§2, 3 are preliminaries for cluster algebras and graded quiver varieties respectively. In §4 we introduce the category $\mathcal{C}_1$ following [31] and study the corresponding graded quiver varieties. In §5 we define a homomorphism from the Grothendieck group $\mathbf{R}_{\ell=1}$ of $\mathcal{C}_1$ to a rational function field which is endowed with a cluster algebra structure. In §6 we explain the relation between the cluster character and the push-forward and derive several consequences on factorizations of simple modules. In §7 we prove that cluster monomials are dual canonical base elements.

Acknowledgments. I began to study cluster algebras after Bernard Leclerc's talk at the meeting 'Enveloping Algebras and Geometric Representation Theory' at the Mathematisches Forschungsinstitut Oberwolfach (MFO) in March 2009. I thank him and David Hernandez for discussions during/after the meeting. They kindly taught me many things on cluster algebras. Alexander Braverman's question/Leclerc's answer (Conway-Coxeter frieze [13]) and discussions with Rina Kedem at the meeting were also very helpful. I thank the organizers, as well as the staffs at the MFO for providing me nice environment for the research. I thank Osamu Iyama for correcting my understanding on the cluster mutation. I thank Andrei Zelevinsky for comments on a preliminary version of this paper. Finally I thank George Lusztig's works which have been a source of my inspiration more than a decade.

2. PRELIMINARIES (I) – CLUSTER ALGEBRAS

We review the definition and properties of cluster algebras.

2.1. Definition. Let $\mathcal{G} = (I, E)$ be a finite graph, where I is the set of vertexes and E is the set of edges. Let H be the set of pairs consisting of an edge together with its orientation. For $h \in H$, we denote by $i(h)$ (resp. $o(h)$) the incoming (resp. outgoing) vertex of h . For $h \in H$ we denote by $\bar{h}$ the same edge as h with the reverse orientation. A *quiver* $\mathcal{Q} = (I, \Omega)$ is the finite graph $\mathcal{G}$ together with a choice of an *orientation* $\Omega \subset H$ such that $\Omega \cap \bar{\Omega} = \emptyset$, $\Omega \cup \bar{\Omega} = H$.

We will consider a pair of a quiver $\mathcal{Q} = (I, \Omega)$ and a larger quiver $\tilde{\mathcal{Q}} = (\tilde{I}, \tilde{\Omega})$ containing $\mathcal{Q}$, where I is a subset of $\tilde{I}$ and Ω is obtained from $\tilde{\Omega}$ by removing arrows incident a point in $\tilde{I} \setminus I$. Set $I_{\text{fr}} = \tilde{I} \setminus I$. We call $i \in I_{\text{fr}}$ (resp. $i \in I$) a *frozen* (resp. *principal*) vertex.

We assume that $\tilde{\mathcal{Q}}$ has no loops nor 2-cycles and there are no edges connecting points in I_{fr} . We define a matrix $\tilde{\mathbf{B}} = (b_{ij})_{i \in \tilde{I}, j \in I}$ by

$$b_{ij} := \begin{aligned} &(\text{the number of oriented edges from } j \text{ to } i) \\ &\text{or } -(\text{the number of oriented edges from } i \text{ to } j). \end{aligned}$$

Since we have assumed $\tilde{\mathcal{Q}}$ contains no 2-cycles, this is well-defined. Moreover, giving $\tilde{\mathbf{B}}$ is equivalent to a quiver $\tilde{\mathcal{Q}}$ with the decomposition $\tilde{I} = I \sqcup I_{\text{fr}}$ as above. The *principal part* $\mathbf{B}$ of $\tilde{\mathbf{B}}$ is the matrix obtained from $\tilde{\mathbf{B}}$ by taking entries for $I \times I$. From the definition $\mathbf{B}$ is skew-symmetric.

For a vertex $k \in I$ we define the *matrix mutation* $\mu_k(\tilde{\mathbf{B}})$ of $\tilde{\mathbf{B}}$ in direction k as the new matrix (b'_{ij}) indexed by $(i, j) \in \tilde{I} \times I$ given by the formula

$$(2.1) \quad b'_{ij} = \begin{cases} -b_{ij} & \text{if } i = k \text{ or } j = k, \\ b_{ij} + \text{sgn}(b_{ik}) \max(b_{ik}b_{kj}, 0) & \text{otherwise.} \end{cases}$$

If $\tilde{\Omega}^*$ denotes the corresponding quiver, it is obtained from $\tilde{\Omega}$ by the following rule:

- (1) For each $i \rightarrow k, k \rightarrow j \in \tilde{\Omega}$, create a new arrow $i \rightarrow j$ if either i or $j \in I$.
- (2) Reverse all arrows incident with k .
- (3) Remove 2-cycles between i and j of the resulting quiver after (1) and (2).

Graphically it is given by

$$\tilde{\Omega} : \begin{array}{ccccc} & & r & & \\ & i & \xrightarrow{\quad} & j & \\ & s \searrow & & \nearrow t & \\ & & k & & \end{array} \quad \Longrightarrow \quad \tilde{\Omega}^* : \begin{array}{ccccc} & & r+st & & \\ & i & \xrightarrow{\quad} & j & \\ & s \searrow & & \nearrow t & \\ & & k & & \end{array}$$

where s, t are nonnegative integers and $i \xrightarrow{l} j$ means that there are l arrows from i to j if $l \geq 0$, $(-l)$ arrows from j to i if $l \leq 0$. The new quiver $\tilde{\Omega}^*$ has no loops nor 2-cycles.

Let $\mathcal{F} = \mathbb{Q}(x_i)_{i \in \tilde{I}}$ be the field of rational functions in commuting indeterminates $\mathbf{x} = (x_i)_{i \in \tilde{I}}$ indexed by $\tilde{I}$. For $k \in I$ we define a new variable x_k^* by the *exchange relation*:

$$(2.2) \quad x_k^* = \frac{\prod_{b_{ik} > 0} x_i^{b_{ik}} + \prod_{b_{ik} < 0} x_i^{-b_{ik}}}{x_k}.$$

Let $\mu_k(\mathbf{x})$ be the set of variables obtained from $\mathbf{x}$ by replacing x_k by x_k^* . The pair $(\mu_k(\mathbf{x}), \mu_k(\tilde{\mathbf{B}}))$ is called the *mutation* of $(\mathbf{x}, \tilde{\mathbf{B}})$ in direction k . We can iterate this procedure and obtain new pairs by mutating $(\mu_k(\mathbf{x}), \mu_k(\tilde{\mathbf{B}}))$ in any direction $l \in I$. We do not make mutations in direction of a frozen vertex $k \in I_{\text{fr}}$. Variables x_i for $i \in I_{\text{fr}}$ are always in $\mu_k(\mathbf{x})$; they are called *frozen variables* (or *coefficients* in [21]).

Now a *seed* is a pair $(\mathbf{y}, \tilde{\mathbf{C}})$ of $\mathbf{y} = (y_i)_{i \in \tilde{I}} \in \mathcal{F}^{\tilde{I}}$ and a matrix $\tilde{\mathbf{C}} = (c_{ij})_{i \in \tilde{I}, j \in I}$ obtained from the *initial seed* $(\mathbf{x}, \tilde{\mathbf{B}})$ by a successive application of mutations in various direction $k \in I$. The set of seeds is denoted by $\mathcal{S}$. A *cluster* is $\{y_i \mid i \in \tilde{I}\}$ of a seed $(\mathbf{y}, \tilde{\mathbf{C}})$, considered as a subset of $\mathcal{F}$ by forgetting the $\tilde{I}$ -index. A *cluster variable* is an element of the union of all clusters. Note that clusters may overlap: a cluster variable may be belonged to another cluster. Also the

$\tilde{I}$ -index may be different from the original one. The *cluster algebra* $\mathcal{A}(\tilde{\mathbf{B}})$ is the subalgebra of $\mathcal{F}$ generated by all the cluster variables. The integer $\#I$ is called the *rank* of $\mathcal{A}(\tilde{\mathbf{B}})$. A *cluster monomial* is a monomial in the cluster variables of a *single* cluster. The exchange relation (2.2) is of the form

$$(2.3) \quad x_k x_k^* = m_+ + m_-$$

where $m_{\pm} = \prod_{\pm b_{ik} > 0} x_i^{\pm b_{ik}}$ are cluster monomials.

When we say a *cluster algebra*, it may mean the subalgebra $\mathcal{A}(\tilde{\mathbf{B}})$ or all the above structures.

One of important results in the cluster algebra theory is the Laurent phenomenon: every cluster variable z in $\mathcal{A}(\tilde{\mathbf{B}})$ is a Laurent polynomial in any given cluster $\mathbf{y}$ with coefficients in $\mathbb{Z}$. It is conjectured that the coefficients are nonnegative. A cluster monomial is a subtraction free rational expression in $\mathbf{x}$, but this is not enough to ensure the positivity of its Laurent expansion, as an example $x^2 - x + 1 = (x + 1)^3 / (x + 1)$ shows.

2.2. F -polynomial. It is known that cluster variables of $\mathcal{A}(\tilde{\mathbf{B}})$ are expressed by the $\mathbf{g}$ -vectors and F -polynomials [23], which are constructed from another cluster algebra with the same principal part, but a simpler frozen part. We recall their definition in this subsection.

We first prepare some notation. We consider the multiplicative group $\mathbb{P}$ of all Laurent monomials in $(x_i)_{i \in I}$. We introduce the addition $\oplus$ by

$$\prod_i x_i^{a_i} \oplus \prod_i x_i^{b_i} = \prod_i x_i^{\min(a_i, b_i)}.$$

This operation together with the ordinary multiplication and division, $\mathbb{P}$ becomes a semifield, called the *tropical semifield*. Let F be a subtraction-free rational expression with integer coefficients in variables y_i . Then we evaluate it in $\mathbb{P}$ by specializing the y_i to some elements p_i of $\mathbb{P}$. We denote it by $F|_{\mathbb{P}}(\mathbf{p})$, where $\mathbf{p} = (p_i)_{i \in I}$.

Let $\mathcal{A}_{\text{pr}}$ be the cluster algebra with *principal coefficients*. It is given by the initial seed $((\mathbf{u}, \mathbf{f}), \tilde{\mathbf{B}}_{\text{pr}})$ with $(\mathbf{u}, \mathbf{f}) = (u_i, f_i)_{i \in I}$, and $\tilde{\mathbf{B}}_{\text{pr}}$ is the matrix indexed by $(I \sqcup I) \times I$ with the same principal $\mathbf{B}$ as $\tilde{\mathbf{B}}$ and the identity matrix in the frozen part. Here I_{fr} is a copy of I and $\tilde{I} = I \sqcup I$. We write a cluster variable α as

$$\alpha = X_{\alpha}(\mathbf{u}, \mathbf{f})$$

a subtraction free rational expression in $\mathbf{u}, \mathbf{f}$. We then specialize all the u_i to 1:

$$F_{\alpha}(\mathbf{f}) = X_{\alpha}(\mathbf{u}, \mathbf{f})|_{u_i=1}.$$

It becomes a polynomial in f_i , and called the F -polynomial ([loc. cit., §3]). It is also known ([loc. cit., §6]) that X_{α} is homogeneous with respect to $\mathbb{Z}^I$ -grading given by

$$\deg u_i = i, \quad \deg f_j = - \sum_i b_{ij} i,$$

where b_{ij} is the matrix entry for the principal part $\mathbf{B}$, and the vertex i is identified with the coordinate vector in $\mathbb{Z}^I$. We then define $\mathbf{g}$ -vector by

$$\mathbf{g}_{\alpha} \stackrel{\text{def.}}{=} \deg X_{\alpha} \in \mathbb{Z}^I.$$

We now return back to the original cluster algebra $\mathcal{A}(\tilde{\mathbf{B}}) \subset \mathbb{Q}(x_i)_{i \in \tilde{I}}$. We introduce the following variables:

$$y_j = \prod_{i \in I_{\text{fr}}} x_i^{b_{ij}}, \quad \hat{y}_j = y_j \prod_{i \in I} x_i^{b_{ij}} \quad (j \in I).$$

We write $\mathbf{y} = (y_i)_{i \in I}$, $\widehat{\mathbf{y}} = (\widehat{y}_i)_{i \in I}$.

We consider the corresponding cluster variable $x[\alpha]$ in the seed of the original cluster algebra $\mathcal{A}(\widetilde{\mathbf{B}})$ obtained by the same mutation processes as we obtained α in the cluster algebra with principal coefficients. We then have [23, Cor. 6.5]:

$$x[\alpha] = \frac{F_\alpha(\widehat{\mathbf{y}})}{F_\alpha|_{\mathbb{P}}(\mathbf{y})} \mathbf{x}^{\mathbf{g}_\alpha},$$

where $\mathbf{x}^{\mathbf{g}_\alpha} = \prod_{i \in I} x_i^{(\mathbf{g}_\alpha)_i}$ if $(\mathbf{g}_\alpha)_i$ is the i^{th} -entry of $\mathbf{g}_\alpha$.

2.3. Hernandez-Leclerc monoidal categorification conjecture. We recall Hernandez-Leclerc's monoidal categorification conjecture in this subsection.

Let $\mathcal{A}$ be a cluster algebra and $\mathcal{M}$ be an abelian monoidal category. A simple object $L \in \mathcal{M}$ is *prime* if there exists no non trivial factorization $L \cong L_1 \otimes L_2$. We say that L is *real* if $L \otimes L$ is simple.

Definition 2.4 ([31]). Let $\mathcal{A}$ and $\mathcal{M}$ as above. We say that $\mathcal{M}$ is a *monoidal categorification* of $\mathcal{A}$ if the Grothendieck ring of $\mathcal{M}$ is isomorphic to $\mathcal{A}$, and if

- (1) the cluster monomials m of $\mathcal{A}$ are the classes of all the real simple objects $L(m)$ of $\mathcal{M}$;
- (2) the cluster variables of $\mathcal{A}$ (including the frozen ones) are the classes of all the real prime simple objects of $\mathcal{M}$.

If two cluster variables x, y belong to the common cluster, then xy is a cluster monomial. Therefore the corresponding simple objects $L(x), L(y)$ satisfy $L(x) \otimes L(y) \cong L(y) \otimes L(x) \cong L(xy)$.

Proposition 2.5 ([31, §2]). *Suppose that a cluster algebra $\mathcal{A}$ has a monoidal categorification $\mathcal{M}$.*

- (1) *Every cluster monomial has a Laurent expansion with positive coefficients with respect to any cluster $\mathbf{y} = (y_i)_{i \in \widetilde{I}} \in \mathcal{S}$;*

$$m = \frac{N_m(\mathbf{y})}{\prod_i y_i^{d_i}}, \quad d_i \in \mathbb{Z}_{\geq 0}, \quad N(y_i) \in \mathbb{Z}_{\geq 0}[y_i^{\pm}].$$

In fact, the coefficient of $\prod y_i^{k_i}$ in $N_m(\mathbf{y})$ is equal to the multiplicity of $L(\prod y_i^{k_i}) = \bigotimes L(y_i)^{\otimes k_i}$ in $L(m) \otimes L(\prod y_i^{d_i}) = L(m) \otimes \bigotimes L(y_i)^{\otimes d_i}$.

- (2) *The cluster monomials of $\mathcal{A}$ are linearly independent.*

Conjecture 2.6 ([31]). *The cluster algebra for the quiver defined in §5 has the monoidal categorification, when the underlying graph is of type ADE. More precisely it is given by a certain monoidal subcategory $\mathcal{C}_1$ of the category of finite dimensional representations of the quantum affine algebra $\mathbf{U}_q(\mathbf{Lg})$.*

The monoidal subcategory will be defined in §4.1 in terms of graded quiver varieties for arbitrary symmetric Kac-Moody cases. And we prove the conjecture for type ADE. This is new for D_n for $n \geq 5$ and E_6, E_7, E_8 since the conjecture was already proved in [31] for type A and D_4 .

However we cannot control the prime factorization of arbitrary simple modules except ADE cases. We can just prove cluster monomials are real simple objects. So it is still not clear that our monoidal subcategory is a monoidal categorification in the above sense for types other than ADE.

3. PRELIMINARIES (II) – GRADED QUIVER VARIETIES

We review the definition of graded quiver varieties and the convolution diagram for the tensor product in this section. Our notation mainly follows [54]. Some materials are borrowed from [58].

We do not explain anything about representations of the quantum loop algebra $\mathbf{U}_q(\mathbf{Lg})$ except in Theorem 3.17. This is because we can work directly in the category of perverse sheaves on graded quiver varieties. Another reason is that it is not known whether the quantum loop algebra $\mathbf{U}_q(\mathbf{Lg})$ can be equipped with the structure of a Hopf algebra in general. Therefore tensor products of modules do not make sense. On the other hand, the category of perverse sheaves *has* the coproduct induced from the convolution diagram.

3.1. Definition of graded quiver varieties. Let q be a nonzero complex number. We will assume that it is not a root of unity later, but can be so at the beginning.

Suppose that a finite graph $\mathcal{G} = (I, E)$ is given. We assume the graph $\mathcal{G}$ contains no edge loops. Let $\mathbf{A} = (a_{ij})$ be the adjacency matrix of the graph, namely

$$a_{ij} = (\text{the number of edges joining } i \text{ to } j).$$

Let $\mathbf{C} = 2\mathbf{I} - \mathbf{A} = (c_{ij})$ be the Cartan matrix.

Let H be the set of pairs consisting of an edge together with its orientation as in §7. We choose and fix a function $\varepsilon: H \rightarrow \mathbb{C}^*$ such that $\varepsilon(h) + \varepsilon(\bar{h}) = 0$ for all $h \in H$. We usually take an orientation Ω of $\mathcal{G}$ and define $\varepsilon(h) = 1$ if $h \in \Omega$ and -1 otherwise.

Let V, W be $I \times \mathbb{C}^*$ -graded vector spaces such that its $(i \times a)$ -component, denoted by $V_i(a)$, is finite dimensional and 0 for all but finitely many $i \times a$. In what follows we consider only $I \times \mathbb{C}^*$ -graded vector spaces with this condition. We say the pair (V, W) of $I \times \mathbb{C}^*$ -graded vector spaces is *l-dominant* if

$$(3.1) \quad \dim W_i(a) - \dim V_i(aq) - \dim V_i(aq^{-1}) - \sum_{j:j \neq i} c_{ij} \dim V_j(a) \geq 0$$

for any i, a .

Let $\mathbf{C}_q$ (q -analog of the Cartan matrix) be an endomorphism of $\mathbb{Z}^{I \times \mathbb{C}^*}$ given by

$$(3.2) \quad (v_i(a)) \mapsto (v'_i(a)); \quad v'_i(a) = v_i(aq) + v_i(aq^{-1}) + \sum_{j:j \neq i} c_{ij} v_j(a).$$

Considering $\dim V, \dim W$ as vectors in $\mathbb{Z}_{\geq 0}^{I \times \mathbb{C}^*}$, we view the left hand side of (3.1) as the (i, a) -component of $(\dim W - \mathbf{C}_q \dim V)$. This is an analog of a weight.

We say $V \leq V'$ if

$$\dim V_i(a) \geq \dim V'_i(a)$$

for any i, a . We say $V < V'$ if $V \leq V'$ and $V \neq V'$. This is analog of the dominance order. We say $(V, W) \leq (V', W')$ if there exists $\mathbf{v}'' \in \mathbb{Z}_{\geq 0}^{I \times \mathbb{C}^*}$ whose entries are 0 for all but finitely many (i, a) such that

$$\dim W - \mathbf{C}_q \dim V = \dim W' - \mathbf{C}_q(\dim V' + \mathbf{v}'').$$

When $W = W'$, $(V, W) \leq (V', W')$ if and only if $V \leq V'$.

These conditions originally come from the representation theory of the quantum loop algebra $\mathbf{U}_q(\mathbf{Lg})$.

For an integer n , we define vector spaces by

$$(3.3) \quad \begin{aligned} L^\bullet(V, W)^{[n]} &\stackrel{\text{def.}}{=} \bigoplus_{i \in I, a \in \mathbb{C}^*} \text{Hom}(V_i(a), W_i(aq^n)), \\ E^\bullet(V, W)^{[n]} &\stackrel{\text{def.}}{=} \bigoplus_{h \in H, a \in \mathbb{C}^*} \text{Hom}(V_{o(h)}(a), W_{i(h)}(aq^n)). \end{aligned}$$

If V and W are $I \times \mathbb{C}^*$ -graded vector spaces as above, we consider the vector spaces

$$(3.4) \quad \mathbf{M}^\bullet \equiv \mathbf{M}^\bullet(V, W) \stackrel{\text{def.}}{=} E^\bullet(V, V)^{[-1]} \oplus L^\bullet(W, V)^{[-1]} \oplus L^\bullet(V, W)^{[-1]},$$

where we use the notation $\mathbf{M}^\bullet$ unless we want to specify V, W . The above three components for an element of $\mathbf{M}^\bullet$ is denoted by B, α, β respectively. (**NB:** In [49] α and β were denoted by i, j respectively.) The $\text{Hom}(V_{o(h)}(a), V_{i(h)}(aq^{-1}))$ -component of B is denoted by $B_{h,a}$. Similarly, we denote by $\alpha_{i,a}, \beta_{i,a}$ the components of α, β .

We define a map $\mu: \mathbf{M}^\bullet \rightarrow L^\bullet(V, V)^{[-2]}$ by

$$\mu_{i,a}(B, \alpha, \beta) = \sum_{i(h)=i} \varepsilon(h) B_{h,aq^{-1}} B_{\bar{h},a} + \alpha_{i,aq^{-1}} \beta_{i,a},$$

where $\mu_{i,a}$ is the (i, a) -component of μ .

Let $G_V \stackrel{\text{def.}}{=} \prod_{i,a} \text{GL}(V_i(a))$. It acts on $\mathbf{M}^\bullet$ by

$$(B, \alpha, \beta) \mapsto g \cdot (B, \alpha, \beta) \stackrel{\text{def.}}{=} (g_{i(h),aq^{-1}} B_{h,a} g_{o(h),a}^{-1}, g_{i,aq^{-1}} \alpha_{i,a}, \beta_{i,a} g_{i,a}^{-1}).$$

The action preserves the subvariety $\mu^{-1}(0)$ in $\mathbf{M}^\bullet$.

Definition 3.5. A point $(B, \alpha, \beta) \in \mu^{-1}(0)$ is said to be *stable* if the following condition holds:
if an $I \times \mathbb{C}^*$ -graded subspace V' of V is B -invariant and contained in $\text{Ker } \beta$, then $V' = 0$.
Let us denote by $\mu^{-1}(0)^s$ the set of stable points.

Clearly, the stability condition is invariant under the action of G_V . Hence we may say an orbit is stable or not.

We consider two kinds of quotient spaces of $\mu^{-1}(0)$:

$$\mathfrak{M}_0^\bullet(V, W) \stackrel{\text{def.}}{=} \mu^{-1}(0) // G_V, \quad \mathfrak{M}^\bullet(V, W) \stackrel{\text{def.}}{=} \mu^{-1}(0)^s / G_V.$$

Here $//$ is the affine algebro-geometric quotient, i.e. the coordinate ring of $\mathfrak{M}_0^\bullet(V, W)$ is the ring of G_V -invariant functions on $\mu^{-1}(0)$. In particular, it is an affine variety. It is the set of closed G_V -orbits. The second one is the set-theoretical quotient, but coincides with a quotient in the geometric invariant theory (see [48, §3]). The action of G_V on $\mu^{-1}(0)^s$ is free thanks to the stability condition ([48, 3.10]). By a general theory, there exists a natural projective morphism

$$\pi: \mathfrak{M}^\bullet(V, W) \rightarrow \mathfrak{M}_0^\bullet(V, W).$$

(See [48, 3.18].) The inverse image of 0 under π is denoted by $\mathfrak{L}^\bullet(V, W)$. We call these varieties *cyclic quiver varieties* or *graded quiver varieties*, according as q is a root of unity or *not*. In this paper we only consider the case q is not a root of unity hereafter. When we want to distinguish $\mathfrak{M}^\bullet(V, W)$ and $\mathfrak{M}_0^\bullet(V, W)$, we call the former (resp. latter) the *nonsingular* (resp. *affine*) *graded quiver variety*. But it does not mean that $\mathfrak{M}_0^\bullet(V, W)$ is actually singular. As we see later, it is possible that $\mathfrak{M}_0^\bullet(V, W)$ happens to be nonsingular.

We have

$$\dim \mathfrak{M}^\bullet(V, W) = \langle \dim V, (q + q^{-1}) \dim W - q^{-1} \mathbf{C}_q \dim V \rangle,$$

where $q^\pm \cdot$ is an automorphism of $\mathbb{Z}^{I \times \mathbb{C}^*}$ given by $(v_i(a)) \mapsto (v'_i(a))$; $v'_i(a) = v_i(aq^\pm)$ and $\langle \cdot, \cdot \rangle$ is the natural pairing on $\mathbb{Z}^{I \times \mathbb{C}^*}$ ([54, 4.11]).

It is known that the coordinate ring of $\mathfrak{M}_0^\bullet(V, W)$ is generated by the following type of elements:

$$(3.6) \quad (B, \alpha, \beta) \mapsto \langle \chi, \beta_{j, aq^{-n-1}} B_{h_n, q^{-n}} \dots B_{h_1, aq^{-1}} \alpha_{i, a} \rangle$$

where χ is a linear form on $\text{Hom}(W_i(a), W_j(aq^{-n-2}))$. (See [44].)

The original quiver varieties [47, 48] are the special case when $q = 1$ and $V_i(a) = W_i(a) = 0$ except $a = 1$. On the other hand, the above varieties $\mathfrak{M}^\bullet(W)$, $\mathfrak{M}_0^\bullet(W)$ are fixed point set of the original quiver varieties with respect to a semisimple element in a product of general linear groups. (See [49, §4].) In particular, it follows that $\mathfrak{M}^\bullet(V, W)$ is nonsingular, since the corresponding original quiver variety is so. This can be also checked directly.

Let $\mathfrak{M}_0^{\bullet \text{reg}}(V, W) \subset \mathfrak{M}_0^\bullet(V, W)$ be a possibly empty open subset of $\mathfrak{M}_0^\bullet(V, W)$ consisting of closed free G_V -orbits. It is known that π is isomorphism on $\pi^{-1}(\mathfrak{M}_0^{\bullet \text{reg}}(V, W))$ [48, 3.24]. In particular, $\mathfrak{M}_0^{\bullet \text{reg}}(V, W)$ is nonsingular and is pure dimensional.

A G_V -orbit though (B, α, β) , considered as a point of $\mathfrak{M}^\bullet(V, W)$ is denoted by $[B, \alpha, \beta]$.

Suppose that we have two $I \times \mathbb{C}^*$ -graded vector spaces V, V' such that $V_i(a) \subset V'_i(a)$ for all i, a . Then $\mathfrak{M}_0^\bullet(V, W)$ can be identified with a closed subvariety of $\mathfrak{M}_0^\bullet(V', W)$ by the extension by 0 to the complementary subspace (see [49, 2.5.3]). We consider the limit

$$\mathfrak{M}_0^\bullet(W) \stackrel{\text{def.}}{=} \bigcup_V \mathfrak{M}_0^\bullet(V, W).$$

(It was denoted by $\mathfrak{M}_0(\infty, \mathbf{w})^A$ in [49], and by $\mathfrak{M}_0^\bullet(\infty, W)$ in [54].)

From the proof of [49, 4.2.2] we have $\mathfrak{M}_0^\bullet(V, 0) = \{0\}$ for $W = 0$ since q is not a root of unity. (The assumption that there is at most one edge joining two vertexes is unnecessary, since our definition of the graded quiver variety is different from one in [loc. cit.] when there are multiple edges joining two vertexes. See Remark 3.13 for more detail.) Then [47, 6.5] or [48, 3.27] implies that

$$(3.7) \quad \mathfrak{M}_0^\bullet(W) = \bigsqcup_{[V]} \mathfrak{M}_0^{\bullet \text{reg}}(V, W),$$

where $[V]$ denotes the isomorphism class of V . It is known that

$$(3.8) \quad \mathfrak{M}_0^{\bullet \text{reg}}(V, W) \neq \emptyset \text{ if and only if } \mathfrak{M}^\bullet(V, W) \neq \emptyset \text{ and } (V, W) \text{ is } l\text{-dominant. (See [49, 14.3.2(2)].)}$$

$$(3.9) \quad \text{If } \mathfrak{M}_0^{\bullet \text{reg}}(V, W) \subset \overline{\mathfrak{M}_0^{\bullet \text{reg}}(V', W)}, \text{ then } V' \leq V. \text{ (This follows from [49, §3.3].)}$$

It is also easy to show that

$$(3.10) \quad \mathfrak{M}_0^{\bullet \text{reg}}(V, W) = \emptyset \text{ if } V \text{ is sufficiently large.}$$

(See the argument in the proof of Proposition 4.6(1).) Thus $\mathfrak{M}_0^\bullet(W) \stackrel{\text{def.}}{=} \bigcup_V \mathfrak{M}_0^\bullet(V, W)$ stabilizes at some V .

On the other hand, we consider the disjoint union for $\mathfrak{M}^\bullet(V, W)$:

$$\mathfrak{M}^\bullet(W) \stackrel{\text{def.}}{=} \bigsqcup_{[V]} \mathfrak{M}^\bullet(V, W).$$

Note that there are no obvious morphisms between $\mathfrak{M}^\bullet(V, W)$ and $\mathfrak{M}^\bullet(V', W)$ since the stability condition is not preserved under the extension. We have a morphism $\mathfrak{M}^\bullet(W) \rightarrow \mathfrak{M}_0^\bullet(W)$, still denoted by π .

It is known that $\mathfrak{M}^\bullet(V, W)$ becomes empty if V is sufficiently large when $\mathfrak{g}$ is of type ADE . (Since the usual quiver variety $\mathfrak{M}(V, W)$ is nonempty if and only if $(\dim W - \mathbf{C} \dim V)$ is a weight of the irreducible representation with the highest weight $\dim W$. See [48, 10.2].) But it is not true in general, and dimensions of $\mathfrak{M}^\bullet(V, W)$ may go to ∞ when V becomes large. In the following, we will use $\mathfrak{M}^\bullet(W)$ as a brevity of the notation, and consider its geometric structure on each $\mathfrak{M}^\bullet(V, W)$ individually. We will never consider it as an infinite dimensional variety. Furthermore, we will only need $\mathfrak{M}^\bullet(V, W)$ such that $\mathfrak{M}_0^{\text{reg}}(V, W) \neq \emptyset$ in practice. From the above remark, we can work stay in finite V 's.

The following three term complex plays an important role:

$$(3.11) \quad C_{i,a}^\bullet(V, W) : V_i(aq) \xrightarrow{\sigma_{i,a}} \bigoplus_{h:i(h)=i} V_{o(h)}(a) \oplus W_i(a) \xrightarrow{\tau_{i,a}} V_i(aq^{-1}),$$

where

$$\sigma_{i,a} = \bigoplus_{i(h)=i} B_{h,aq} \oplus \beta_{i,aq}, \quad \tau_{i,a} = \sum_{i(h)=i} \varepsilon(h) B_{h,a} + \alpha_{i,a}.$$

This is a complex thanks to the equation $\mu(B, \alpha, \beta) = 0$. If (B, α, β) is stable, $\sigma_{i,a}$ is injective as an $I \times \mathbb{C}^*$ -graded vector space V' given by $V'_i(a) := \text{Ker } \sigma_{i,a}$, $V'_j(b) := 0$ (otherwise) is B -invariant and contained in $\text{Ker } \beta$, and hence must be 0.

We assign the degree 0 to the middle term. We define the rank of complex $C^\bullet$ by $\sum_p (-1)^p \text{rank } C^p$. It is exactly the left hand side of (3.1). Therefore (V, W) is l -dominant if and only if

$$\text{rank } C_{i,a}^\bullet(V, W) \geq 0$$

for any i, a . From this observation the ‘only-if’ part of (3.8) is clear: If we consider the complex at a point $\mathfrak{M}_0^{\text{reg}}(V, W)$, it is easy to see $\tau_{i,a}$ is surjective. Therefore $\text{rank } C_{i,a}^\bullet(V, W)$ is the dimension of the middle cohomology group. When (V, W) is l -dominant, we define an $I \times \mathbb{C}^*$ -graded vector space $C^\bullet(V, W)$ by

$$(3.12) \quad \dim (C^\bullet(V, W))_i(a) = \text{rank } C_{i,a}^\bullet(V, W).$$

Remark 3.13. Since we only treat graded quiver varieties of type ADE in [54], we explain what must be modified for general types.

In [49] the graded quiver varieties are the $\mathbb{C}^*$ -fixed points of the ordinary quiver varieties. When there are multiple edges joining two vertexes, there are several choices of the $\mathbb{C}^*$ -action. A choice corresponds to a choice of the q -analog $\mathbf{C}_q$ of the Cartan matrix $\mathbf{C}$ which implicitly appears in the defining relation of the quantum loop algebras. See [loc. cit., (1.2.9)] for the defining relation and [loc. cit., (2.9.1)] or (3.11) for its relation to the $\mathbb{C}^*$ -action. For example, consider type $A_1^{(1)}$. In [loc. cit.] the q -analog of the Cartan matrix was

$$\begin{pmatrix} [2]_q & [-2]_q \\ [-2]_q & [2]_q \end{pmatrix} = \begin{pmatrix} q + q^{-1} & -(q + q^{-1}) \\ -(q + q^{-1}) & q + q^{-1} \end{pmatrix},$$

while it is

$$\begin{pmatrix} [2]_q & -2 \\ -2 & [2]_q \end{pmatrix} = \begin{pmatrix} q + q^{-1} & -2 \\ -2 & q + q^{-1} \end{pmatrix}$$

in this paper. When there is at most one edge joining two vertexes, we do not have these choices as $[1]_q = 1$. The theory developed in [49] works for any choice of the $\mathbb{C}^*$ -action.

For results in [54], we need a little care. First of all, [loc. cit., Cor. 3.7] does not make sense since it is not known whether we have tensor products in general as we already mentioned. For the choice of the $\mathbb{C}^*$ -action in this paper, all other results hold without any essential changes,

except assertions when ε is a root of unity or ± 1 . (In these cases, we will get new types of strata so the assertion must be modified. For the affine type, they can be understood from [52].) If we take the $\mathbb{C}^*$ -action in [49], the recursion used to prove **Axiom 2** does not work. So we first take the $\mathbb{C}^*$ -action in this paper, and then apply the same trick used to deal with cyclic quiver varieties. In particular, we need to include analog of **Axiom 4**. Details are left as an exercise for the reader of [54].

3.2. Transversal slice. Take a point $x \in \mathfrak{M}_0^{\bullet \text{reg}}(V^0, W)$. Let T be the tangent space of $\mathfrak{M}_0^{\bullet \text{reg}}(V^0, W)$ at x . Since $\mathfrak{M}_0^{\bullet \text{reg}}(V^0, W)$ is nonempty, (V^0, W) is l -dominant, i.e. (3.1) holds by (3.8). Let $W^\perp = C^\bullet(V^0, W)$ as in (3.12).

We consider another graded quiver variety $\mathfrak{M}_0^\bullet(V, W)$ which contains x in its closure. By (3.9) we have $V \leq V^0$. Therefore we can consider $V^\perp$, $I \times \mathbb{C}^*$ -graded vector space whose (i, a) -component has the dimension $\dim V_i(a) - \dim V_i^0(a)$. We have $\dim W - \mathbf{C}_q \dim V = \dim W^\perp - \mathbf{C}_q \dim V^\perp$, which means the ‘weight’ is unchanged under this procedure.

Theorem 3.14 ([49, §3.3]). *We work in the complex analytic topology. There exist neighborhoods U , U_T , $U_\mathfrak{S}$ of $x \in \mathfrak{M}_0^\bullet(V, W)$, $0 \in T$, $0 \in \mathfrak{M}_0^\bullet(V^\perp, W^\perp)$ respectively, and biholomorphic maps $U \rightarrow U_T \times U_\mathfrak{S}$, $\pi^{-1}(U) \rightarrow U_T \times \pi^{-1}(U_\mathfrak{S})$ such that the following diagram commutes:*

$$\begin{array}{ccc} \mathfrak{M}^\bullet(V, W) \supset \pi^{-1}(U) & \xrightarrow[\cong]{} & U_T \times \pi^{-1}(U_\mathfrak{S}) \subset T \times \mathfrak{M}_0^\bullet(V^\perp, W^\perp) \\ \pi \downarrow & & \downarrow \text{id} \times \pi \\ \mathfrak{M}_0^\bullet(V, W) \supset U & \xrightarrow[\cong]{} & U_T \times U_\mathfrak{S} \subset T \times \mathfrak{M}_0^\bullet(V^\perp, W^\perp) \end{array}$$

Furthermore, a stratum $\mathfrak{M}_0^{\bullet \text{reg}}(V', W)$ of $\mathfrak{M}_0^\bullet(V, W)$ is mapped to a product of U_T and the stratum $\mathfrak{M}_0^{\bullet \text{reg}}(V'^\perp, W^\perp)$ of $\mathfrak{M}_0^\bullet(V^\perp, W^\perp)$.

Here $V'^\perp$ is defined exactly as $V^\perp$ replacing V by V' , i.e. $\dim V'^\perp = \dim V' - \dim V^0$.

Note that $V'' \leq V' \Leftrightarrow V''^\perp \leq V'^\perp$ if we define $V''^\perp$ for V'' in the same way.

See also [14] for the same result in the étale topology.

3.3. The additive category $\mathcal{Q}_W$ and the Grothendieck ring. Let X be a complex algebraic variety. Let $\mathcal{D}(X)$ be the bounded derived category of constructible sheaves of $\mathbb{C}$ -vector spaces on X . For $j \in \mathbb{Z}$, the shift functor is denoted by $L \mapsto L[j]$. The Verdier duality is denoted by D . For a locally closed subvariety $Y \subset X$, we denote by 1_Y the constant sheaf on Y . We denote by $IC(Y)$ the intersection cohomology complex associated with the trivial local system 1_Y on Y . Our degree convention is so that $IC(Y)|_Y = 1_Y[\dim Y]$.

Since $\pi: \mathfrak{M}^\bullet(V, W) \rightarrow \mathfrak{M}_0^\bullet(V, W)$ is proper and $\mathfrak{M}^\bullet(V, W)$ is smooth, $\pi_!(1_{\mathfrak{M}^\bullet(V, W)})$ is a direct sum of shifts of simple perverse sheaves on $\mathfrak{M}_0^\bullet(V, W)$ by the decomposition theorem [2]. We denote by $\mathcal{P}_W$ the set of isomorphism classes of simple perverse sheaves obtained in this manner, considered as a complex on $\mathfrak{M}_0^\bullet(W)$ by extension by 0 to the complement of $\mathfrak{M}_0^\bullet(V, W)$. By [49, §14] $\mathcal{P}_W = \{IC(\mathfrak{M}_0^{\bullet \text{reg}}(V, W)) \mid \mathfrak{M}_0^{\bullet \text{reg}}(V, W) \neq \emptyset\}$. By (3.10) $\#\mathcal{P}_W < \infty$. Set $IC_W(V) \stackrel{\text{def.}}{=} IC(\mathfrak{M}_0^{\bullet \text{reg}}(V, W))$. Let $\mathcal{Q}_W$ be the full subcategory of $\mathcal{D}(\mathfrak{M}_0^\bullet(W))$ whose objects are the complexes isomorphic to finite direct sums of $IC_W(V)[k]$ for various $IC_W(V) \in \mathcal{P}_W$, $k \in \mathbb{Z}$. Let $\pi_W(V) \stackrel{\text{def.}}{=} \pi_!(1_{\mathfrak{M}^\bullet(V, W)}[\dim \mathfrak{M}^\bullet(V, W)])$. By the definition, we have $\pi_W(V) \in \mathcal{Q}_W$. The subcategory $\mathcal{Q}_W$ is preserved under D and elements in $\mathcal{P}_W$ are fixed by D .

Let $\mathcal{K}(\mathcal{Q}_W)$ be the abelian group with one generator (L) for each isomorphism class of objects of $\mathcal{Q}_W$ and with relations $(L) + (L') = (L'')$ whenever L'' is isomorphic to $L \oplus L'$. It is a module over $\mathcal{A} = \mathbb{Z}[t, t^{-1}]$ by $t(L) = (L[1])$, $t^{-1}(L) = (L[-1])$. It is a free $\mathcal{A}$ -module with

base $\{(IC_W(V)) \mid IC_W(V) \in \mathcal{P}_W\}$. The duality D defines the *bar involution* $\overline{}$ on $\mathcal{K}(\mathcal{Q}_W)$ fixing $(IC_W(V))$ and satisfying $\overline{t(L)} = t^{-1}(\overline{L})$. Since π is proper and $\mathfrak{M}^\bullet(V, W)$ is smooth, we also have $\overline{(\pi_W(V))} = (\pi_W(V))$. We do not write () hereafter.

There is another base

$$\{\pi_W(V) \mid (V, W) \text{ is } l\text{-dominant, } \mathfrak{M}^\bullet(V, W) \neq \emptyset\}.$$

Note that $\pi_W(V)$ make sense for any V without the l -dominance condition, but we need to take only l -dominant ones to have a base. Let us define $a_{V,V';W}(t)$ by

$$(3.15) \quad \pi_W(V) = \sum_{V'} a_{V,V';W}(t) IC_W(V').$$

Then we have $a_{V,V';W}(t) \in \mathbb{Z}_{\geq 0}[t, t^{-1}]$, $a_{V,V;W}(t) = 1$ and $a_{V,V';W} = 0$ unless $V' \leq V$. Since both $\pi_W(V)$ and $IC_W(V')$ are fixed by the bar involution, we have $a_{V,V';W}(t) = a_{V,V';W}(t^{-1})$. It also follows that we only need to consider $\pi_!(1_{\mathfrak{M}^\bullet(V,W)})$ for which (V, W) is l -dominant in the definition of $\mathcal{P}_W$.

Take V^0 such that $\mathfrak{M}_0^{\text{reg}}(V^0, W) \neq \emptyset$. Taking account the transversal slice in §3.2, we define a surjective homomorphism $p_{W^\perp, W}: \mathcal{K}(\mathcal{Q}_W) \rightarrow \mathcal{K}(\mathcal{Q}_{W^\perp})$ by

$$IC_W(V) \mapsto \begin{cases} IC_{W^\perp}(V^\perp) & \text{if } \mathfrak{M}_0^{\text{reg}}(V^0, W) \subset \overline{\mathfrak{M}_0^{\text{reg}}(V, W)}, \\ 0 & \text{otherwise.} \end{cases}$$

By Theorem 3.14 this homomorphism is also compatible with $\pi_W(V)$. Taking various V 's, $\mathcal{K}(\mathcal{Q}_W)$'s form a projective system.

We consider the dual $\mathcal{K}(\mathcal{Q}_W)^* = \text{Hom}_{\mathcal{A}}(\mathcal{K}(\mathcal{Q}_W), \mathcal{A})$. Let $\{L_W(V)\}, \{\chi_W(V)\}$ be the bases of dual to $\{IC_W(V)\}, \{\pi_W(V)\}$ respectively. Here V runs over the set of isomorphism classes of $I \times \mathbb{C}^*$ -graded vector spaces such that (V, W) is l -dominant. We consider yet another base $\{M_W(V)\}$ of $\mathcal{K}(\mathcal{Q}_W)^*$ given by

$$\mathcal{K}(\mathcal{Q}_W) \ni (L) \mapsto \sum_k t^{\dim \mathfrak{M}_0^{\text{reg}}(V, W) - k} \dim H^k(i_{x_{V,W}}^! L) \in \mathcal{A},$$

where $x_{V,W}$ is a point in $\mathfrak{M}_0^{\text{reg}}(V, W)$ and $i_{x_{V,W}}$ is the inclusion of the point $x_{V,W}$ in $\mathfrak{M}_0^\bullet(W)$. By Theorem 3.14 it is independent of the choice of $x_{V,W}$. Also it is compatible with the projective system: if $V^0 \geq V' \geq V$, $\langle M_W(V'), IC_W(V) \rangle = \langle M_{W^\perp}(V'^\perp), IC_{W^\perp}(V^\perp) \rangle$.

By the defining property of perverse sheaves, we have

$$(3.16) \quad L_W(V) \in M_W(V) + \sum_{V': V' > V} t^{-1} \mathbb{Z}[t^{-1}] M_W(V').$$

Since there are only finitely many V' with $V' > V$, this is a finite sum. This shows that $\{M_W(V)\}_V$ is a base. Recall also that the canonical base $L_W(V)$ is characterized by this property together with $\overline{L_W(V)} = L_W(V)$. It is the analog of the characterization of Kazhdan-Lusztig base. This is not relevant in this paper, but was important to compute $L_W(V)$ explicitly in [54].

Let

$$\mathbf{R}_t \stackrel{\text{def.}}{=} \left\{ (f_W) \in \prod_W \text{Hom}_{\mathcal{A}}(\mathcal{K}(\mathcal{Q}_W), \mathcal{A}) \mid \begin{array}{l} \langle f_W, x_W \rangle = \langle f_{W^\perp}, p_{W^\perp, W}(x_W) \rangle \\ \text{for any } W, W^\perp, x_W \in \mathcal{K}(\mathcal{Q}_W) \end{array} \right\}.$$

A functional $(f_W) \in \mathbf{R}_t$ is determined when all values $\langle f_{W^\perp}, IC_{W^\perp}(0) \rangle$ are given for any $W^\perp$. Let $L(W)$, $\chi(W)$, $M(W)$ be the functional determined from $L_W(0)$, $\chi_W(0)$, $M_W(0)$

respectively. For example, $\langle L(W), IC_{W'}(V') \rangle = \delta_{\dim W, \dim W' - \mathbf{C}_q \dim V'}$. They form analog of *canonical*, *monomial* and *PBW* bases of $\mathbf{R}_t$ respectively. From (3.16) the transition matrix between the canonical and monomial bases are upper triangular with respect to the ordering $(0, W) \leq (0, W')$.

The following is the main result in [49].

Theorem 3.17 ([49, 14.3.10]). *As an abelian group, $\mathbf{R}_t|_{t=1}$ is isomorphic to the Grothendieck group of the category $\mathcal{R}$ of l -integrable representations of the quantum loop algebra $\mathbf{U}_q(\mathbf{Lg})$ of the symmetric Kac-Moody Lie algebra $\mathbf{g}$ given by the Cartan matrix $\mathbf{C}$ so that*

- $L(W)$ corresponds to the class of the simple module whose Drinfeld polynomial is given by

$$P_i(u) = \prod_{a \in \mathbb{C}^*} (1 - au)^{\dim W_i(a)} \quad (i \in I).$$

- $M(W)$ corresponds to the class of the standard module whose Drinfeld polynomial is given by the same formula.

Since we do not need this result in this paper, except for an explanation of our approach to one in [31], we do not explain terminologies and concepts in the statement. See [49].

From a general theory of the convolution algebra (see [12]), $\mathcal{K}(\mathcal{Q}_W)$ is the Grothendieck group of the category of *graded* representations of the convolution algebra $H_*(\mathfrak{M}^\bullet(W) \times_{\mathfrak{m}_0(W)} \mathfrak{M}^\bullet(W)) \cong \bigoplus_{V^1, V^2} \text{Ext}_{\mathcal{Q}(\mathfrak{m}_0(W))}^*(\pi_W(V^1), \pi_W(V^2))$, where the grading is for $\text{Ext}^\bullet$ -group. And $\{L_W(V)\}$ is the base given by classes of simple modules.

Let us briefly explain how we glue the abelian categories for various W to get a single abelian category. A family of graded module structures $\{\rho_W: H_*(\mathfrak{M}^\bullet(W) \times_{\mathfrak{m}_0(W)} \mathfrak{M}^\bullet(W)) \rightarrow \text{End}_{\mathbb{C}}(V)\}_W$ on a single vector space M is said to be *compatible*, if ρ_W factors through various restrictions to open subsets in Theorem 3.14 and the restrictions are compatible with the restriction of $\rho_{W^\perp}$ under the local isomorphisms in Theorem 3.14. For example, we fix W^0 and choose various points $x_{V,W} \in \mathfrak{m}_0^{\text{reg}}(V, W)$ with $\dim W - \mathbf{C}_q \dim V = \dim W^0$. We identify $H_*(\pi^{-1}(x_{V,W}))$ with a single vector space M , say $H_*(\pi^{-1}(x_{0,W^0}))$, by the local isomorphisms. It is a compatible family of module structures. Compatible families form an abelian category. Let us denote it by $\mathcal{R}_{\text{conv}}$. Then we have $\mathcal{K}(\mathcal{R}_{\text{conv}}) \cong \mathbf{R}_t$. In the above theorem, we have families of homomorphisms $\mathbf{U}_q(\mathbf{Lg}) \rightarrow H_*(\mathfrak{M}^\bullet(W) \times_{\mathfrak{m}_0(W)} \mathfrak{M}^\bullet(W))$ compatible with the local isomorphisms. Therefore we have a functor from $\mathcal{R}_{\text{conv}}$ to the category $\mathcal{R}$ of l -integrable representations of $\mathbf{U}_q(\mathbf{Lg})$. It sends a simple object to a simple module. We do not know whether it is an equivalence (after forgetting the grading on $\mathcal{R}_{\text{conv}}$), but we can get enough information practically.

3.4. t -analog of q -characters. For each $(i, a) \in I \times \mathbb{C}^*$, we introduce an indeterminate $Y_{i,a}$. Let

$$\mathcal{Y}_t \stackrel{\text{def.}}{=} \mathcal{A}[Y_{i,a}, Y_{i,a}^{-1}]_{i \in I, a \in \mathbb{C}^*}.$$

We associate polynomials $e^W, e^V \in \mathcal{Y}_t$ to graded vector spaces V, W by

$$e^W = \prod_{i \in I, a \in \mathbb{C}^*} Y_{i,a}^{\dim W_i(a)}, \quad e^V = \prod_{i \in I, a \in \mathbb{C}^*} V_{i,a}^{\dim V_i(a)}, \quad \text{where } V_{i,a} = Y_{i,aq^{-1}}^{-1} Y_{i,aq}^{-1} \prod_{\substack{h \in H \\ o(h)=i}} Y_{i(h),a}.$$

We define the t -analog of q -character for $M(W)$ by

$$\chi_{q,t}(M(W)) \stackrel{\text{def.}}{=} \sum_V \sum_k t^{-k} \dim H^k(i_0^! \pi_W(V)) e^W e^V,$$

where 0 is the unique point of $\mathfrak{M}_0^\bullet(0, W)$. From the definition in the previous subsection, this is nothing but the generating function of pairings $\langle M_W(0), \pi_W(V) \rangle$ for various V . If $\mathfrak{g}$ is of type ADE , $\mathfrak{M}^\bullet(V, W)$ becomes empty for large V . (Since the usual quiver variety $\mathfrak{M}(V, W)$ is nonempty if and only if $(\dim W - \mathbf{C} \dim V)$ is a weight of the irreducible representation with the highest weight $\dim W$. See [48, 10.2].) Therefore this is a finite sum. If $\mathfrak{g}$ is not of type ADE , this becomes an infinite series, so it lives in a completion of $\mathscr{Y}_t$. Since the difference is not essential, we keep the notation $\mathscr{Y}_t$. Anyway we will only use the truncated q -character in this paper.

Suppose V^0 is l -dominant and we define $V^\perp, W^\perp$ as in §3.2. Then

$$\sum_V \langle M_W(V^0), \pi_W(V) \rangle = \sum_V \langle M_{W^\perp}(0), \pi_{W^\perp}(V^\perp) \rangle = \chi_{q,t}(M(W^\perp))$$

as $e^W e^V = e^{W^\perp} e^{V^\perp}$.

Since $\{M(W)\}$ is a base of $\mathbf{R}_t$, we can extend $\chi_{q,t}$ to $\mathbf{R}_t$ linearly. We have

$$(3.18) \quad \chi_{q,t}(L(W)) = \sum_V \langle L_W(0), \pi_W(V) \rangle e^W e^V = \sum_V a_{V,0;W}(t) e^W e^V,$$

where $a_{V,0;W}$ is the coefficient of $IC_W(0) = 1_{\{0\}}$ in $\pi_W(V)$ (in $\mathcal{K}(\mathscr{Q}_W)$) as in (3.15).

Since $\{M_W(V)\}_{(V,W):l\text{-dominant}}$ forms a base of $\mathbf{R}_t$, we have

Theorem 3.19. *The q -character homomorphism $\chi_{q,t}: \mathbf{R}_t \rightarrow \mathscr{Y}_t$ is injective.*

But $\chi_{q,t}$ also contains terms from $\pi_W(V)$ with (V, W) not necessary l -dominant. These are redundant information.

Remark 3.20. By [54, Th. 3.5] the coefficient of $e^W e^V$ in the t -analog of q -characters for standard modules $M(W)$ is in $t^{\dim \mathfrak{M}^\bullet(V,W)} \mathbb{Z}_{\geq 0}[t^{-2}]$. This was a consequence of vanishing of odd cohomology groups of $\mathfrak{L}^\bullet(V, W)$. From the proof of [12, Lem. 8.7.8] together with the above vanishing result, we have

$$a_{V,0;W}(t) \in t^{\dim \mathfrak{M}^\bullet(V,W)} \mathbb{Z}_{\geq 0}[t^{-2}].$$

3.5. A convolution diagram. Let us take a 2-step flag $0 \subset W^2 \subset W$ of $I \times \mathbb{C}^*$ -graded vector spaces. We put $W/W^2 = W^1$. Following [50], we introduce closed subvarieties in $\mathfrak{M}_0^\bullet(W)$ and $\mathfrak{M}^\bullet(W)$:

$$\begin{aligned} \mathfrak{Z}_0^\bullet(W^1; W^2) &= \{[B, \alpha, \beta] \in \mathfrak{M}_0^\bullet(W) \mid W^2 \text{ is invariant under } \beta B^k \alpha \text{ for any } k \in \mathbb{Z}_{\geq 0}\}, \\ \mathfrak{Z}^\bullet(W^1; W^2) &= \pi^{-1}(\mathfrak{Z}_0^\bullet(W^1; W^2)). \end{aligned}$$

This definition is different from the original one, but equivalent [loc. cit., 3.6, 3.7]. The latter has an α -partition

$$\mathfrak{Z}^\bullet(W^1; W^2) = \bigsqcup \mathfrak{Z}^\bullet(V^1, W^1; V^2, W^2)$$

such that $\mathfrak{Z}^\bullet(V^1, W^1; V^2, W^2)$ is a vector bundle over $\mathfrak{M}^\bullet(V^1, W^1) \times \mathfrak{M}^\bullet(V^2, W^2)$ of rank

$$\langle \dim V^1, q^{-1}(\dim W^2 - \mathbf{C}_q \dim V^2) \rangle + \langle \dim V^2, q \dim W^1 \rangle.$$

(See [loc. cit., 3.8].) Let us denote this rank by

$$d(V^1, W^1; V^2, W^2).$$

(It was denote by $d(e^{V^1} e^{W^1}, e^{V^2} e^{W^2})$ in [54].)

Following [58] we consider the diagram

$$\mathfrak{M}_0^\bullet(W^1) \times \mathfrak{M}_0^\bullet(W^2) \xleftarrow{\kappa} \mathfrak{Z}_0^\bullet(W^1; W^2) \xrightarrow{\iota} \mathfrak{M}_0^\bullet(W),$$

where ι is the inclusion and κ is given by the induced maps from $\beta B^k \alpha$ to $W^1 = W/W^2$, W^2 . Then we define a functor

$$\tilde{\text{Res}}_{W^1, W^2} \stackrel{\text{def.}}{=} \kappa_! \iota^*: \mathcal{D}(\mathfrak{M}_0^\bullet(W)) \rightarrow \mathcal{D}(\mathfrak{M}_0^\bullet(W^1) \times \mathfrak{M}_0^\bullet(W^2)).$$

We have

$$\tilde{\text{Res}}_{W^1, W^2}(\pi_W(V)) = \bigoplus_{V^1+V^2=V} \pi_{W^1}(V^1) \boxtimes \pi_{W^2}(V^2) [d(V^2, W^2; V^1, W^1) - d(V^1, W^1; V^2, W^2)].$$

(See [58, Lemma 4.1]. A weaker statement was given in [54, 6.2(3)].) From this observation objects in $\mathcal{Q}_W$ are sent to $\mathcal{Q}_{W^1 \times W^2}$, the full subcategory of $\mathcal{D}(\mathfrak{M}_0^\bullet(W^1) \times \mathfrak{M}_0^\bullet(W^2))$ whose objects are complexes isomorphic to finite direct sums of $IC_{W^1}(V^1) \boxtimes IC_{W^2}(V^2)[k]$ for various $IC_{W^1}(V^1) \in \mathcal{P}_{W^1}$, $IC_{W^2}(V^2) \in \mathcal{P}_{W^2}$, $k \in \mathbb{Z}$ ([58, Lemma 4.1]). Therefore this functor induces a homomorphism $\mathcal{K}(\mathcal{Q}_W) \rightarrow \mathcal{K}(\mathcal{Q}_{W^1}) \otimes_{\mathcal{A}} \mathcal{K}(\mathcal{Q}_{W^2})$. It also shows that it is coassociative, as $\mathcal{K}(\mathcal{Q}_W)$ is spanned by classes $\tilde{\pi}_W(V)$ and they satisfy the coassociativity from the above formula. We denote it also by $\tilde{\text{Res}}_{W^1, W^2}$.

Let $\mathbf{C}_q^{-1}$ be the inverse of $\mathbf{C}_q$. We define it by solving the equation $(u_i(a)) = \mathbf{C}_q(x_i(a))$ recursively starting from $x_i(aq^s) = 0$ for sufficiently small s . Note that $x_i(a)$ may be nonzero for infinitely many a . We then observe

$$d(V^1, W^1; V^2, W^2) - \langle \mathbf{C}_q^{-1} \dim W^1, q^{-1} \dim W^2 \rangle$$

is preserved under the replacement $\mathfrak{M}^\bullet(V^1, W^1) \times \mathfrak{M}^\bullet(V^2, W^2) \rightsquigarrow \mathfrak{M}^\bullet(V^{1\perp}, W^{1\perp}) \times \mathfrak{M}^\bullet(V^{2\perp}, W^{2\perp})$ by the transversal slice ([58, Lemma 3.2]). Therefore we define

$$\begin{aligned} \varepsilon(W^1, W^2) &\stackrel{\text{def.}}{=} \langle \mathbf{C}_q^{-1} \dim W^1, q^{-1} \dim W^2 \rangle - \langle \mathbf{C}_q^{-1} \dim W^2, q^{-1} \dim W^1 \rangle, \\ \text{Res} &\stackrel{\text{def.}}{=} \sum_{W=W^1 \oplus W^2} \tilde{\text{Res}}[\varepsilon(W^1, W^2)] \end{aligned}$$

Then its transpose defines a multiplication on $\mathbf{R}_t$, which is denoted by $\otimes$.

We also define the twisted multiplication on $\mathcal{Y}_t$ given by

$$(3.21) \quad m_1 * m_2 = t^{\varepsilon(\vec{m}_1, \vec{m}_2)} m_1 m_2,$$

where m_1, m_2 are monomials in $Y_{i,a}^\pm$ and $\vec{m}_\alpha = (m_i^\alpha(a))$ is given by $m_\alpha = \prod Y_{i,a}^{m_i^\alpha(a)}$.

The following is the main result of [58].

Theorem 3.22. (1) *The structure constant of the product with respect to the base $\{L(W)\}$ is positive:*

$$L(W^1) \otimes L(W^2) \in \sum_W a_{W^1, W^2}^W(t) L(W)$$

with $a_{W^1, W^2}^W(t) \in \mathbb{Z}_{\geq 0}[t, t^{-1}]$.

(2) $\chi_{q,t}: \mathbf{R}_t \rightarrow \mathcal{Y}_t$ is an algebra homomorphism with respect to $\otimes$ and the twisted product $*$.

The following corollary of the positivity is also due to [58].

Corollary 3.23. *The followings are equivalent:*

- (1) $L(W^1) \otimes L(W^2) = L(W^1 \oplus W^2)$ holds at $t = 1$.
- (2) $L(W^1) \otimes L(W^2) = t^{\varepsilon(W^1, W^2)} L(W^1 \oplus W^2)$.

It is tiresome to keep powers of t when tensor products of simple modules are simple. From this corollary, there is no loss of information even if we forget powers. Therefore we do not write $t^{\varepsilon(W^1, W^2)}$ hereafter.

The restriction functor defines an algebra homomorphism

$$H_*(\mathfrak{M}^\bullet(W) \times_{\mathfrak{M}_0^\bullet(W)} \mathfrak{M}^\bullet(W)) \rightarrow H_*(\mathfrak{M}^\bullet(W^1) \times_{\mathfrak{M}_0^\bullet(W^1)} \mathfrak{M}^\bullet(W^1)) \otimes H_*(\mathfrak{M}^\bullet(W^2) \times_{\mathfrak{M}_0^\bullet(W^2)} \mathfrak{M}^\bullet(W^2)).$$

It gives us a monoidal structure on the un-graded version of $\mathcal{R}_{\text{conv}}$.

4. GRADED QUIVER VARIETIES FOR THE MONOIDAL SUBCATEGORY $\mathcal{C}_1$

4.1. Graded quiver varieties and the decorated quiver. The monoidal subcategory $\mathcal{C}_1$ introduced in [31] is, in fact, the first (or second) of series of subcategories $\mathcal{C}_\ell$ indexed by $\ell \in \mathbb{Z}_{\geq 0}$. Let us describe all of them in terms of the category $\mathcal{R}_{\text{conv}}$.

We suppose that (I, E) contains no odd cycles and take a *bipartite* partition $I = I_0 \sqcup I_1$, i.e. every edge connects a vertex in I_0 with one in I_1 . We set

$$\xi_i = \begin{cases} 0 & \text{if } i \in I_0, \\ 1 & \text{if } i \in I_1. \end{cases}$$

Fix a nonnegative integer ℓ . We consider the graded quiver varieties $\mathfrak{M}^\bullet(V, W)$, $\mathfrak{M}_0^\bullet(V, W)$ under the following condition

$$(*_\ell) \quad W_i(a) = 0 \text{ unless } a = q^{\xi_i}, q^{\xi_i+2}, \dots, q^{\xi_i+2\ell}.$$

It is clear that if W satisfies $(*_\ell)$, both W^1 and W^2 satisfy $(*_\ell)$ in the convolution product $\text{Res}: \mathcal{Q}_W \rightarrow \mathcal{Q}_{W^1} \times \mathcal{Q}_{W^2}$. Also from the proof of Proposition 4.6(1) below, it is clear that $\mathfrak{M}_0^{\text{reg}}(V, W) \neq \emptyset$ implies $V_i(a) = 0$ unless $a = q^{\xi_i+1}, \dots, q^{\xi_i+2\ell-1}$. Since $W_i^\perp(a)$ in §3.2 is the middle cohomology of the complex (3.11), $W_i^\perp(a)$ also satisfies $(*_\ell)$. Therefore the condition $(*_\ell)$ is also compatible with the projective system $\mathcal{K}(\mathcal{Q}_W) \rightarrow \mathcal{K}(\mathcal{Q}_{W^\perp})$. Therefore we have the subring $\mathbf{R}_{t,\ell}$ of $\mathbf{R}_t$. We set $\mathbf{R}_\ell = \mathbf{R}_{t,\ell}|_{t=1}$. It is also clear that the definition in [31] in terms of roots of Drinfeld polynomials corresponds to our definition when $\mathfrak{g}$ is of type *ADE* from the theory developed in [49].

Example 4.1. Consider the simplest case $\ell = 0$. By [49, 4.2.2] or the argument below we have $\mathfrak{M}_0^\bullet(V, W) = \{0\}$ if W satisfies $(*_0)$. Therefore $\mathcal{Q}_W$ consists of finite direct sums of shifts of a single object $1_{\mathfrak{M}_0^\bullet(0,W)}$. We have $\text{Res}(1_{\mathfrak{M}_0^\bullet(0,W)}) = 1_{\mathfrak{M}_0^\bullet(0,W^1)} \boxtimes 1_{\mathfrak{M}_0^\bullet(0,W^2)}$. This corresponds to the fact that any tensor product of simple modules in $\mathcal{C}_0$ remains simple. (See [31, 3.3].)

We now start to analyze the condition $(*_{\ell=1})$. Let

$$(4.2) \quad \mathbf{E}_W \stackrel{\text{def.}}{=} \bigoplus_i \text{Hom}(W_i(q^{\xi_i+2}), W_i(q^{\xi_i})) \oplus \bigoplus_{h: o(h) \in I_1, i(h) \in I_0} \text{Hom}(W_{o(h)}(q^3), W_{i(h)}(1))$$

This vector space $\mathbf{E}_W$ is the space of representations of the *decorated quiver*.

Definition 4.3. Suppose that a finite graph $\mathcal{G} = (I, E)$ together with a bipartite partition $I = I_0 \sqcup I_1$ is given. We define the *decorated quiver* $\tilde{\mathcal{Q}} = (\tilde{I}, \tilde{\Omega})$ by the following two steps.

(1) We put an orientation to each edge in E so that vertexes in I_0 (resp. I_1) are sinks (resp. sources). Let Ω be the set of all oriented edges and $\mathcal{Q} = (I, \Omega)$ be the corresponding quiver.

(2) Let I_{fr} be a copy of I . For $i \in I$, we denote by i' the corresponding vertex in I_{fr} . Then we add a new vertex i' and an arrow $i' \rightarrow i$ (resp. $i \rightarrow i'$) if $i \in I_0$ (resp. $i \in I_1$) for each $i \in I$. Let Ω_{dec} be the set of these arrows. The decorated quiver is $\tilde{\mathcal{Q}} = (\tilde{I}, \tilde{\Omega}_{\text{dec}}) = (I \sqcup I_{\text{fr}}, \Omega \sqcup \Omega_{\text{dec}})$.

We call $\mathcal{Q} = (I, \Omega)$ the *principal part* of the decorated quiver.

For example, for type A_3 with $I_0 = \{1, 3\}$, we get the following quiver:

$$(4.4) \quad \begin{array}{ccccc} W_1(1) & \xleftarrow{\mathbf{y}_{1,2}=\beta_{1,q}B_{1,2,q^2}\alpha_{2,q^3}} & W_2(q^3) & \xrightarrow{\mathbf{y}_{3,2}=\beta_{3,q}B_{3,2,q^2}\alpha_{2,q^3}} & W_3(1) \\ \uparrow \mathbf{x}_1=\beta_{1,q}\alpha_{1,q^2} & & \downarrow \mathbf{x}_2=\beta_{2,q^2}\alpha_{2,q^3} & & \uparrow \mathbf{x}_3=\beta_{3,q}\alpha_{3,q^2} \\ W_1(q^2) & & W_2(q) & & W_3(q^2) \end{array}$$

The maps attached with arrows will soon be clear in the proof of Proposition 4.6.

The following is a variant of a variety corresponding to a monomial in F_i in Lusztig's theory [43, 9.1.3].

Definition 4.5. (1) Let $\nu = (\nu_i) \in \mathbb{Z}_{\geq 0}^I$. Let $\mathcal{F}(\nu, W)$ be the variety parametrizing collections of vector spaces $X = (X_i)_{i \in I}$ indexed by I such that $\dim X_i = \nu_i$ and

$$X_i \subset W_i(1) \ (i \in I_0), \quad X_i \subset W_i(q) \oplus \bigoplus_{h \in \Omega: o(h)=i} X_{i(h)} \ (i \in I_1)$$

It is a kind of a partial flag variety and nonsingular projective.

(2) Let $\tilde{\mathcal{F}}(\nu, W)$ be the variety of all triples $(\bigoplus \mathbf{x}_i, \bigoplus \mathbf{y}_h, X)$ where $(\bigoplus \mathbf{x}_i, \bigoplus \mathbf{y}_h) \in \mathbf{E}_W$ and $X \in \mathcal{F}(\nu, W)$ such that

$$\text{Im } \mathbf{x}_i \subset X_i \ (i \in I_0), \quad \text{Im } \left(\mathbf{x}_i \oplus \bigoplus_{h \in \Omega: o(h)=i} \mathbf{y}_h \right) \subset X_i \ (i \in I_1).$$

Let $\pi_\nu: \tilde{\mathcal{F}}(\nu, W) \rightarrow \mathbf{E}_W$ be the natural projection.

Proposition 4.6. Suppose W satisfies $(*_\ell)$ with $\ell = 1$.

(1) If $\mathfrak{M}_0^{\bullet \text{reg}}(V, W) \neq \emptyset$, we have

$$(4.7) \quad V_i(a) = 0 \text{ unless } a = q^{\xi_i+1}.$$

Moreover we have an isomorphism $\mathfrak{M}_0^\bullet(W) \cong \mathbf{E}_W$ given by

$$[B, \alpha, \beta] \mapsto \left(\bigoplus_{i \in I} \mathbf{x}_i, \bigoplus_{h \in \Omega} \mathbf{y}_h \right); \quad \mathbf{x}_i = \beta_{i,q^{\xi_i+1}} \alpha_{i,q^{\xi_i+2}}, \quad \mathbf{y}_h = \beta_{i(h),q} B_{h,q^2} \alpha_{o(h),q^3}.$$

(2) Suppose that V satisfies (4.7). Let us define $\nu \in \mathbb{Z}_{\geq 0}^I$ by $\nu_i = \dim V_i(q^{\xi_i+1})$. Then $\mathfrak{M}^\bullet(V, W)$ is isomorphic to $\tilde{\mathcal{F}}(\nu, W)$ and the following diagram is commutative:

$$\begin{array}{ccc} \mathfrak{M}^\bullet(V, W) & \xrightarrow{\cong} & \tilde{\mathcal{F}}(\nu, W) \\ \pi \downarrow & & \downarrow \pi_\nu \\ \mathfrak{M}_0^\bullet(W) & \xrightarrow{\cong} & \mathbf{E}_W \end{array}$$

Proof. (1) Consider a map

$$\beta_{j,aq^{-n-1}} B_{h_n,q^{-n}} \dots B_{h_1,aq^{-1}} \alpha_{i,a}: W_i(a) \rightarrow W_j(aq^{-n-2})$$

with $i(h_a) = o(h_{a+1})$ for $a = 1, \dots, n-1$. From the assumption $(*)_1$, this is nonzero only when $i = j$, $n = 0$, $a = q^{\xi_i+2}$ or $n = 1$, $i \in I_1$, $j \in I_0$, $a = q^3$. From this observation we have

$$\mathfrak{M}_0^\bullet(W) = \mathfrak{M}_0^\bullet(V, W),$$

for some V with $V_i(a) = 0$ unless $a = q^{\xi_i+1}$. Thus we obtain the first assertion. Moreover, the equation $\mu(B, \alpha, \beta) = 0$ is automatically satisfied, and the second assertion follows from a standard fact $\text{Hom}(W, V) \oplus \text{Hom}(V, W') // \text{GL}(V) \cong \text{Hom}(W, W')$ for V with $\dim V \geq \min(\dim W, \dim W')$.

(2) We first observe the following:

Claim. Under the assumption (B, α, β) is stable if and only if the following linear maps are all injective:

$$\beta_{i,q}: V_i(q) \rightarrow W_i(1) \ (i \in I_0), \quad \sigma_{i,q}: V_i(q^2) \rightarrow \bigoplus_{h:o(h)=i} V_{i(h)}(q) \oplus W_i(q) \ (i \in I_1).$$

(See (3.11) and the subsequent formula for the definition of $\sigma_{i,q}$.)

Consider the $I \times \mathbb{C}^*$ -graded vector space given by $V'_i(q) = \text{Ker } \beta_{i,q}$ and all other $V'_j(a) = 0$. Then the stability condition implies $V'_i(q) = 0$. Therefore $\beta_{i,q}$ is injective. The same argument shows the injectivity of $\sigma_{i,q}$. Conversely suppose all the above maps are injective. If V' is an $I \times \mathbb{C}^*$ -graded subspace of V as in the definition of the stability condition. (See Definition 3.5.) First consider $V'_i(q)$ for $i \in I_0$. We have $\beta_{i,q}|V'_i(q) = 0$. Therefore the injectivity of $\beta_{i,q}$ implies $V'_i(q) = 0$. Next consider $V'_j(q^2) \subset V_j(q^2)$ for $j \in I_0$. We have $\beta_{j,q^2}|V'_j(q^2) = 0$ from the assumption. We also have $B_{h,q^2}(V'_j(q^2)) \subset V'_i(q) = 0$ from what we have just proved. Therefore the injectivity of $\sigma_{j,q}$ implies that $V'_j(q^2) = 0$. This completes the proof of the claim.

Suppose $[B, \alpha, \beta] \in \mathfrak{M}^\bullet(V, W)$ is given. We set

$$\begin{aligned} \tilde{\sigma}_{i,q} &:= \left(\bigoplus_{h:o(h)=i} \beta_{i(h),q} \oplus \text{id}_{W_i(q)} \right) \circ \sigma_{i,q}: V_i(q^2) \rightarrow \bigoplus_{h:o(h)=i} W_{i(h)}(1) \oplus W_i(q), \\ X_i &:= \text{Im } \beta_{i,q} \ (i \in I_0), \quad X_i := \text{Im } \tilde{\sigma}_{i,q} \ (i \in I_1). \end{aligned}$$

The spaces X_i are independent of the choice of a representative (B, α, β) of $[B, \alpha, \beta]$. From the above claim, we have $\dim X_i = \dim V_i(q)$ ($i \in I_0$) and $\dim X_i = \dim V_i(q^2)$ ($i \in I_1$). The remaining properties are automatically satisfied by the construction.

Conversely suppose that $(\bigoplus \mathbf{x}_i, \bigoplus \mathbf{y}_h, X)$ is given. We set $V_i(q) := X_i$ ($i \in I_0$), $V_i(q^2) := X_i$ ($i \in I_1$) and define linear maps (B, α, β) by

$$\begin{aligned} \beta_{i,q} &:= (\text{the inclusion } X_i \subset W_i(1)), \quad \alpha_{i,q^2} := \mathbf{x}_i \ (i \in I_0), \\ \beta_{i,q^2} \oplus \bigoplus_{h:o(h)=i} B_{h,q^2} &:= \left(\text{the inclusion } X_i \subset W_i(q) \oplus \bigoplus_{h:o(h)=i} X_{i(h)} \right), \quad \alpha_{i,q^3} := \mathbf{x}_i \ (i \in I_1). \end{aligned}$$

From the claim, the data (B, α, β) is stable and defines a point in $\mathfrak{M}^\bullet(V, W)$. These two assignments are inverse to each other, hence they are isomorphisms. $\square$

4.2. A contravariant functor σ . For a later application we study the description in Proposition 4.6(2) further. By (2) $\mathfrak{M}^\bullet(V, W) \cong \tilde{\mathcal{F}}(\nu, W)$ can be considered as a vector bundle over $\mathcal{F}(\nu, W)$. It is naturally a subbundle of the trivial bundle $\mathcal{F}(\nu, W) \times \mathbf{E}_W$. Let $\tilde{\mathcal{F}}(\nu, W)^\perp$ be

its annihilator in the dual trivial bundle $\mathcal{F}(V, W) \times \mathbf{E}_W^*$ and let $\pi^\perp: \tilde{\mathcal{F}}(\nu, W)^\perp \rightarrow \mathbf{E}_W^*$ be the natural projection. We denote the dual variables of $\mathbf{x}_i, \mathbf{y}_h$ by $\mathbf{x}_i^*, \mathbf{y}_h^*$ respectively, i.e.

$$\mathbf{x}_i^* \in \text{Hom}(W_i(q^{\xi_i}), W_i(q^{\xi_i+2})), \quad \mathbf{y}_h^* \in \text{Hom}(W_{i(h)}(1), W_{o(h)}(q^3)).$$

By (2) $(\bigoplus \mathbf{x}_i^*, \bigoplus \mathbf{y}_h^*)$ is contained in $\tilde{\mathcal{F}}(\nu, W)^\perp$ if and only if

$$\mathbf{x}_i^*(X_i) = 0 \ (i \in I_0), \quad \left(\mathbf{x}_i^* + \sum_{h:o(h)=i} \mathbf{y}_h^* \right) (X_i) = 0 \ (i \in I_1).$$

It will be important to understand a fiber of $\pi^\perp$ on a general point $(\bigoplus \mathbf{x}_i^*, \bigoplus \mathbf{y}_h^*)$ in $\mathbf{E}_W^*$. Since considering a subspace X_i in $W_i(q) \oplus W_{i(h)}(1)$ looks slightly strange, let us apply the Bernstein-Gelfand-Ponomarev reflection functors [5] (see [1, VII.5]) to $(\bigoplus \mathbf{x}_i^*, \bigoplus \mathbf{y}_h^*)$ at all the vertexes $i \in I_1$ (where $W_i(q^3)$ is put). First observe that $(\pi^\perp)^{-1}(\bigoplus \mathbf{x}_i^*, \bigoplus \mathbf{y}_h^*)$ is unchanged even if we replace $W_i(q^3)$ by the image of the map

$$(4.8) \quad \mathbf{x}_i^* + \sum_{h:o(h)=i} \mathbf{y}_h^*: W_i(q) \oplus \bigoplus_{h:o(h)=i} W_{i(h)}(1) \rightarrow W_i(q^3)$$

for all $i \in I_1$. Then we set

$$\sigma W_i(q^3) \stackrel{\text{def.}}{=} \text{Ker} \left(\mathbf{x}_i^* + \sum_{h:o(h)=i} \mathbf{y}_h^* \right),$$

and define linear maps $\sigma \mathbf{x}_i: \sigma W_i(q^3) \rightarrow W_i(q)$ ($i \in I_1$), $\sigma \mathbf{y}_h: \sigma W_i(q^3) \rightarrow W_{i(h)}(1)$ ($h \in H$ with $o(h) = i \in I_1$) as the compositions of the inclusion $\sigma W_i(q^3) \rightarrow W_i(q) \oplus \bigoplus_{h:o(h)=i} W_{i(h)}(1)$ and the projections to factors. We have

$$(4.9) \quad \dim \sigma W_i(q^3) = \max \left(\dim W_i(q) + \sum_{h:o(h)=i} \dim W_{i(h)}(1) - \dim W_i(q^3), 0 \right).$$

We denote by σW the new $I \times \mathbb{C}^*$ -graded vector space given obtained from W by replacing $W_i(q^3)$ by $\sigma W_i(q^3)$ for all $i \in I_1$. We also set $\sigma \mathbf{x}_i = \mathbf{x}_i^*$ for $i \in I_0$. We do not change $W_i(1)$, $W_i(q^2)$ for $i \in I_0$ and $W_i(q)$ for $i \in I_1$.

Lemma 4.10. *Let $\sigma \mathbf{x}_i, \sigma \mathbf{y}_h$ be as above. Then $(\pi^\perp)^{-1}(\bigoplus \mathbf{x}_i^*, \bigoplus \mathbf{y}_h^*)$ is isomorphic to the variety of $I \times \mathbb{C}^*$ -graded subspaces X of σW satisfying*

$$\begin{aligned} X_i(q^2) &= 0 \ (i \in I_0), & X_i(q) &= \sigma W_i(q) \ (i \in I_1), \\ \dim X_i(1) &= \dim V_i(q) \ (i \in I_0), & \dim X_i(q^3) &= \dim V_i(q^2) \ (i \in I_1), \\ X &\text{ is invariant under } \left(\bigoplus_{i \in I} \sigma \mathbf{x}_i, \bigoplus_{h \in \Omega} \sigma \mathbf{y}_h \right). \end{aligned}$$

This variety is what people call the *quiver Grassmannian* associated with the quiver representation $(\bigoplus_i \sigma \mathbf{x}_i, \bigoplus_{h \in \Omega} \sigma \mathbf{y}_h)$. Its importance in the cluster algebra theory was first noticed by [7]. We will be interested only in its Poincaré polynomial, which is independent of the choice of a general point, we denote this variety simply by $\text{Gr}_V(\sigma W)$, suppressing the choice $(\bigoplus_i \sigma \mathbf{x}_i, \bigoplus_{h \in \Omega} \sigma \mathbf{y}_h)$. Note also that the I -grading is only relevant in $\text{Gr}_V(\sigma W)$. Therefore we use this notation also for an I -graded vector space V .

Note that the orientation is different from the decorated quiver (4.4). This corresponds to the cluster algebra with principal coefficients considered in §2.2. Therefore we call it the quiver with *principal decoration*. For example, in type A_3 with $I_0 = \{1, 3\}$, we get the following quiver:

$$(4.11) \quad \begin{array}{ccccc} W_1(1) & \xleftarrow{\sigma_{\mathbf{y}_{1,2}}} & \sigma W_2(q^3) & \xrightarrow{\sigma_{\mathbf{y}_{3,2}}} & W_3(1) \\ \sigma_{\mathbf{x}_1=\mathbf{x}_1^*} \downarrow & & \downarrow \sigma_{\mathbf{x}_2} & & \downarrow \sigma_{\mathbf{x}_3=\mathbf{x}_3^*} \\ W_1(q^2) & & W_2(q) & & W_3(q^2) \end{array}$$

Remark 4.12. The quiver Grassmannian is a fiber of a projective morphism, which played a fundamental role in Lusztig's construction of the canonical base. It is denoted by $\pi_{\mathbf{v}}: \tilde{\mathcal{F}}_{\mathbf{v}} \rightarrow \mathbf{E}_{\mathbf{v}}$ in [43, Part II]. But note that Lusztig considered more generally various spaces of *flags* not only subspaces.

Later it will be useful to view σ as a functor between category of representations of quivers. Let $\text{rep} \tilde{\mathcal{Q}}$ be the category of finite dimensional representations of the decorated quiver $\tilde{\mathcal{Q}}$. Let ${}^{\sigma} \tilde{\mathcal{Q}}$ be the quiver with the principal decoration obtained by reversing the arrows between i and i' for $i \in I_0$ as above. Let $\text{rep} {}^{\sigma} \tilde{\mathcal{Q}}$ be the corresponding category and $\text{rep} {}^{\sigma} \tilde{\mathcal{Q}}^{\text{op}}$ be its opposite category. Then σ is the functor

$$\sigma(\bullet) = \prod_{i \in I_1} \Phi_i^- \circ D(\bullet): \text{rep} \tilde{\mathcal{Q}} \rightarrow \text{rep} {}^{\sigma} \tilde{\mathcal{Q}}^{\text{op}},$$

where Φ_i^- is the reflection functor at the vertex for $W_i(q^3)$ and D is the duality operator

$$D(\bullet) = \text{Hom}_{\mathbb{C}}(\bullet, \mathbb{C}).$$

In order to make an identification with the above picture, we fix an isomorphism $W \cong W^*$ of $(I \sqcup I_{\text{fr}})$ -graded vector spaces.

Let $\text{rep}^- \tilde{\mathcal{Q}}$ be the full subcategory of $\text{rep} \tilde{\mathcal{Q}}$ consisting of representations having no direct summand isomorphic to simple modules corresponding to vertexes $i \in I_1$. Similarly we define $\text{rep}^- {}^{\sigma} \tilde{\mathcal{Q}}^{\text{op}}$. Then σ defines an equivalence between $\text{rep}^- \tilde{\mathcal{Q}}$ and $\text{rep}^- {}^{\sigma} \tilde{\mathcal{Q}}^{\text{op}}$. We write the quasi-inverse functor $\sigma_- = D \circ \prod_{i \in I_1} \Phi_i^+$.

In fact, it is more elegant to consider σ as a functor between derived categories of $\text{rep} \tilde{\mathcal{Q}}$ and $\text{rep} {}^{\sigma} \tilde{\mathcal{Q}}^{\text{op}}$ as in [27, IV.4.Ex. 6]. See also Remark 7.7.

5. FROM GROTHENDIECK RINGS TO CLUSTER ALGEBRAS

Since W always satisfies $(*_\ell=1)$ hereafter, we denote $W_i(q^{3\xi_i})$ and $W_i(q^{2-\xi_i})$ by W_i and $W_{i'}$ respectively. This is compatible with the notation in Definition 4.3 as $W_i(q^{2-\xi_i})$ is on the new vertex i' .

We denote the simple modules of the decorated quiver by $S_i, S_{i'}$ corresponding to vertexes $i \in I, i' \in I_{\text{fr}}$. We will consider modules of two completely different algebras, (a) modules in $\mathcal{R}_{\text{conv}}$ (or of $\mathbf{U}_q(\mathbf{Lg})$) and (b) modules of the decorated quiver. Simple modules for the former will be denoted by $L(W)$, while $S_i, S_{i'}$ for the latter. We hope there will be no confusion. We denote the underlying $\tilde{I} = (I \sqcup I_{\text{fr}})$ -graded vector space of $S_i, S_{i'}$ also by the same letter.

The Grothendieck ring $\mathbf{R}_{\ell}$ is a polynomial ring in the class of the classes $L(W)$ with $\dim W = 1$ satisfying $(*_\ell)$ (l -fundamental representations in $\mathcal{C}_{\ell}$ when $\mathbf{g}$ is of type ADE). This result was proved as a consequence of the theory of q -characters in [31, Prop. 3.2] for $\mathbf{g}$ of type ADE .

Since q -characters make sense for arbitrary $\mathfrak{g}$, the same argument works. The corresponding result for the whole category $\mathcal{R}$ is well-known.

For $\mathbf{R}_{\ell=1}$, we have $2\#I$ variables corresponding to l -fundamental representations. We denote them by x_i and x'_i exchanging i and i' from the index of the decorated quiver (Definition 4.3):

$$(5.1) \quad x_i = L(W) \longleftrightarrow W = S_{i'}, \quad x'_{i'} = L(W) \longleftrightarrow W = S_i.$$

This is confusing, but we cannot avoid it to get a correct statement.

We denote the class of the Kirillov-Reshetikhin module in $\mathcal{C}_1$ by f_i . It corresponds to the class $L(W)$, where W is a 2-dimensional $\tilde{I}$ -graded vector space with $\dim W_i = \dim W_{i'} = 1$, and 0 at other gradings. We have

$$(5.2) \quad f_i = x_i x'_{i'} - \prod_{h \in H: o(h)=i} x_{i(h)}.$$

This is an example of the T -system proved in [53], but in fact, easy to check by studying the convolution diagram as $\mathbf{E}_W \cong \mathbb{C}$ has only two strata, the origin and the complement. It is also a simple consequence of Theorem 6.3 below. It is a good exercise for the reader.

Remark 5.3. In [53] more precise relation was shown in the level of modules, not only in the Grothendieck group: for $i \in I_0$, there exists a short exact sequence

$$0 \rightarrow \bigotimes_{h \in H: o(h)=i} x_{i(h)} \rightarrow x'_{i'} \otimes x_i \rightarrow f_i \rightarrow 0$$

and we replace the middle term by $x_i \otimes x'_{i'}$ if $i \in I_1$.

We have an algebra embedding

$$\mathbf{R}_{\ell=1} = \mathbb{Z}[x_i, x'_{i'}]_{i \in I} \rightarrow \mathcal{F} = \mathbb{Q}(x_i, f_i)_{i \in I}.$$

We now put the cluster algebra structure on the right hand side. It is enough to specify the initial seed. We take x_i, f_i as cluster variables of the initial seed. We make f_i as a frozen variable. We call the quiver for the initial seed the $\mathbf{x}$ -quiver. It looks almost the same as the decorated quiver in Definition 4.3, but a little different.

Definition 5.4. Suppose that a finite graph $\mathcal{G} = (I, E)$ together with a bipartite partition $I = I_0 \sqcup I_1$ is given. We define the $\mathbf{x}$ -quiver $\tilde{\mathcal{Q}}_{\mathbf{x}} = (\tilde{I}, \tilde{\Omega}_{\mathbf{x}})$ by the following two steps.

(1) The underlying graph is the same as one of the decorated quiver: $\mathcal{G} = (I \sqcup I_{\text{fr}}, E \sqcup \{i - i'\})$. The variable x_i corresponds to the vertex i in the original quiver, while f_i corresponds to the new vertex i' .

(2) The rule for drawing arrows is

$$(5.5) \quad f_i \rightarrow x_i \ (i \in I_0), \quad x_i \rightarrow f_i \ (i \in I_1), \quad x_{o(h)} \xrightarrow{h} x_{i(h)} \ (\text{if } o(h) \in I_0, i(h) \in I_1).$$

For our favorite example, A_3 with $I_0 = \{1, 3\}$, we get the following quiver.

$$\begin{array}{ccccc} x_1 & \longrightarrow & x_2 & \longleftarrow & x_3 \\ \uparrow & & \downarrow & & \uparrow \\ f_1 & & f_2 & & f_3 \end{array}$$

Note that the orientation differs from the decorated quiver (4.4) nor the principal decoration (4.11). Also the vertex f_i corresponds to $W_{i'}$, and x_i corresponds to W_i . This is different from the identification (5.1). If we look at the principal part, the orientation is reversed.

If we make a mutation in direction x_i , the new variable given by the exchange relation (2.2) is nothing but

$$x'_i = \frac{f_i + \prod_{h \in H: o(h)=i} x_{i(h)}}{x_i}$$

from (5.2). Note the exchange relation is correct for the $\mathbf{x}$ -quiver given by our rule (5.5), but *wrong* for the decorated quiver. Thus this confusion cannot be avoided.

We thus have

Proposition 5.6. *The Grothendieck ring $\mathbf{R}_{\ell=1}$ is a subalgebra of the cluster algebra $\mathcal{A}(\tilde{\mathbf{B}})$.*

The argument in [31, 4.4] (based on [3, 1.21]) implies that $\mathbf{R}_{\ell=1} \cong \mathcal{A}(\tilde{\mathbf{B}})$, but we will see that all cluster monomials come from simple modules in $\mathbf{R}_{\ell=1}$, so we have a different proof later.

We also need the seed obtained by applying the sequence of mutations $\prod_{i \in I_1} \mu_i$. Then (1) x_i ($i \in I_1$) is replaced by x'_i , (2) the orientation of arrows are reversed in the principal part and $i \rightarrow i'$ ($i \in I_1$), and (3) add a_{ij} arrows from i to j' . In our A_3 example, we obtain

$$(5.7) \quad \begin{array}{ccccc} & x_1 & \longleftarrow & x'_2 & \longrightarrow & x_3 \\ & \uparrow & & \uparrow & & \uparrow \\ f_1 & & \searrow & & \swarrow & \\ & f_2 & & & & f_3 \end{array}$$

We set

$$(5.8) \quad z_i \stackrel{\text{def.}}{=} \begin{cases} x_i & \text{if } i \in I_0, \\ x'_i & \text{if } i \in I_1. \end{cases}$$

We call above one the $\mathbf{z}$ -quiver.

6. CLUSTER CHARACTER AND PRIME FACTORIZATIONS OF SIMPLE MODULES

6.1. An almost simple module. Fix an $\tilde{I}$ -graded vector space W . Let Ψ be the Fourier-Sato-Deligne functor for the vector space $\mathbf{E}_W \cong \mathfrak{M}_0^*(W)$ ([36, 39]). We define a subset $\mathcal{L}_W \subset \mathcal{P}_W$ by

$$L \in \mathcal{L}_W \iff \text{the support of } \Psi(L) \text{ is the whole space } \mathbf{E}_W^*.$$

If $L \in \mathcal{L}_W$, $\Psi(L)$ is an IC complex associated with a local system defined over an open set in $\mathbf{E}_W$. We denote its rank by $r_W(L) \in \mathbb{Z}_{>0}$.

Since the Fourier transform of $IC_W(0) = 1_{\{0\}}$ is $1_{\mathbf{E}_W^*}[\dim \mathbf{E}_W^*]$, we always have $IC_W(0) \in \mathcal{L}_W$. We have $r_W(IC_W(0)) = 1$.

We extend this definition for a condition on simple modules $L(W')$. Recall that $IC_W(V)$ is identified with $IC_{W^\perp}(0)$ such that $\dim W^\perp = \dim W - \mathbf{C}_q \dim V$. We say $L(W') \in \mathcal{L}_W$ if $IC_W(V) \in \mathcal{L}_W$ with $W' = W^\perp$. We similarly define $r_W(L(W'))$.

We define the *almost simple module* associated with W by

$$\mathbb{L}(W) = \sum_{L(W') \in \mathcal{L}_W} r_W(L(W')) L(W').$$

This is an element in $\mathbf{R}_t$.

From the definition of $L(W') \in \mathcal{L}_W$ we have $W' \leq W$. Therefore almost simple modules $\{\mathbb{L}(W)\}$ form a basis of $\mathbf{R}_t$ such that the transition matrix between it and $\{L(W)\}$ is upper triangular with diagonal entries 1.

We will see that an almost simple module is not necessarily simple later. There will be also a simple sufficient condition guaranteeing an almost simple module is simple.

Remark 6.1. As we will see soon, almost simple modules are given in terms of quiver Grassmannian for a generic representation of $\mathbf{E}_W^*$. This, at first sight, looks similar to the set of generic variables considered by Dupont [20]. (See also [19].) But there is a crucial difference. We consider the total sum of Betti numbers of the quiver Grassmannian, while Dupont consider Euler numbers. There is an example with nontrivial odd degree cohomology groups [17, Ex. 3.5], so this is really difference.

Note that from the representation theory of $\mathbf{U}_q(\mathbf{Lg})$, it is natural to specialize as $t = 1$, since t -analog becomes the ordinary q -character (and the positivity is preserved). This difference cannot be seen for cluster monomials, thanks to Remark 3.20.

6.2. Truncated q -character. In [31, §6] Hernandez-Leclerc introduced the *truncated q -character* $\chi_q(M)_{\leq 2}$ from the ordinary q -character $\chi_q(M)$ by setting variables $V_{i,q^r} = 0$ for $r \geq 3$. From the geometric definition of the q -character reviewed in §3.4, it just means that we only consider nonsingular quiver varieties $\mathfrak{M}^\bullet(V, W)$ satisfying (4.7), i.e. those studied in Proposition 4.6(2). In particular, its t -analog also makes sense:

$$(6.2) \quad \begin{aligned} \chi_{q,t}(M(W))_{\leq 2} &\stackrel{\text{def.}}{=} \sum_{V \text{ satisfies (4.7)}} \sum_k t^{-k} \dim H^k(i_0^! \pi_W(V)) e^W e^V, \\ \chi_{q,t}(L(W))_{\leq 2} &= \sum_{V \text{ satisfies (4.7)}} a_{V,0;W}(t) e^W e^V, \end{aligned}$$

where $a_{V,0;W}(t)$ is the coefficient of $IC_W(0) = 1_{\{0\}}$ in $\pi_W(V)$ in $\mathcal{K}(\mathcal{Q}_W)$. Since V satisfies (4.7) if (V, W) is l -dominant, the truncated q -character still embed $\mathbf{R}_{\ell=1}$ to $\mathcal{U}_t$. (See [31, Prop. 6.1] for an algebraic proof.)

The following is one of main results in this paper.

Theorem 6.3. *Suppose W satisfies $(*_\ell)$ with $\ell = 1$. Then the truncated t -analog of q -character of an almost simple module is given by*

$$\chi_{q,t}(\mathbb{L}(W))_{\leq 2} = \sum_V P_t(\text{Gr}_V(\sigma W)) e^W e^V,$$

where the summation runs over all $I \times \mathbb{C}^*$ -graded vector spaces V with (4.7) and $P_t(\)$ is the normalized Poincaré polynomial for the Borel-Moore homology group

$$P_t(\text{Gr}_V(\sigma W)) = \sum_i t^{i - \dim \mathfrak{M}^\bullet(V, W)} \dim H_i(\text{Gr}_V(\sigma W)).$$

Since $\text{Gr}_V(\sigma W)$ is a fiber of $\pi^\perp: \tilde{\mathcal{F}}(\nu, W)^\perp \rightarrow \mathbf{E}_W^*$ over a general point in $\mathbf{E}_W^*$ and $\tilde{\mathcal{F}}(\nu, W)^\perp$ is nonsingular, $\text{Gr}_V(\sigma W)$ is nonsingular by the generic smoothness theorem. Since $\pi^\perp$ is projective, it is also projective. Therefore the Poincaré polynomial is essentially equal to the virtual one defined by Danilov-Khovanskii [15] using a mixed Hodge structure of Deligne [16]:

$$P_t^{\text{vert}}(X) \stackrel{\text{def.}}{=} \sum_k (-1)^k t^{p+q} h^{p,q}(H_c^k(X)).$$

(See [15] for the notation $h^{p,q}(H_c^k(X))$.) Since our Poincaré polynomial is normalized, we have

$$P_t(\text{Gr}_V(\sigma W)) = t^{-\dim \mathfrak{M}_0^{\bullet, \text{reg}}(V, W)} P_{-t}^{\text{vert}}(\text{Gr}_V(\sigma W)).$$

Remark 6.4. Recall that $\chi_{q,t}(L(W))$ was computed in [54]. More precisely, a purely combinatorial algorithm to compute $\chi_{q,t}(L(W))$ was given in [loc. cit.]. If we are interested in simple modules in $\mathcal{C}_1$, the same algorithm works by replacing every ' $\chi_{q,t}(\cdot)$ ' in [loc. cit.] by ' $\chi_{q,t}(\cdot)_{\leq 2}$ '. Thus the computation is drastically simplified. The algorithm consists of 3 steps. The first step is the computation of $\chi_{q,t}$ for l -fundamental representation. The actual computation of $\chi_{q,t}$ was performed by a supercomputer [55]. But this is certainly unnecessary for $\chi_{q,t}(\cdot)_{\leq 2}$. The second step is the computation of $\chi_{q,t}$ for the standard modules. This is just a twisted multiplication of $\chi_{q,t}$'s given in the first step. This step is simple. The third step is analog of the definition of Kazhdan-Lusztig polynomials. It is still hard computation if we take large W . It is probably interesting to compare this algorithm with one given by the mutation, e.g., for W corresponding to the highest root of E_8 . In this case we have $\mathbb{L}(W) = L(W)$ as we will see soon in Proposition 6.9.

In general, if $\mathbb{L}(W) \neq L(W)$, we need to compute $r_W(L(W'))$.

Example 6.5. For the Kirillov-Reshetikhin module f_i , we have $\dim W_i = 1 = \dim W_{i'}$. If $i \in I_1$, we have $\sigma W = 0$. Therefore $\text{Gr}_V(\sigma W)$ is a point if $V = 0$ and $\emptyset$ otherwise. If $i \in I_0$, a generic $\sigma \mathbf{x}_i^*: W_i \rightarrow W_{i'}$ is an isomorphism. Therefore $\text{Gr}_V(\sigma W)$ is again a point if $V = 0$ and $\emptyset$ otherwise. Thus we must have $\mathbb{L}(W) = L(W)$ in this case, and $\chi_{q,t}(f_i)_{\leq 2}$ contains only the first term:

$$\chi_{q,t}(f_i)_{\leq 2} = Y_{i,q^{\xi_i}} Y_{i,q^{\xi_i+2}}.$$

This can be shown in many ways, say using the main result of [53].

Next consider x_i . If $i \in I_0$, then σW is 1-dimensional with nonzero entry at $\sigma W_{i'}$. But since we can put only 0-dimensional space $X_{i'}$, we only allow $V = 0$. Thus $\mathbb{L}(W) = L(W)$ and $\chi_{q,t}(x_i) = Y_{i,q^2}$.

If $i \in I_1$, then σW is 2-dimensional with nonzero entries at σW_i and $\sigma W_{i'}$. Therefore we have either $V = 0$ or 1-dimensional V with nonzero entry at $V_{i'}$. The corresponding varieties are a single point in both cases. Thus $\mathbb{L}(W) = L(W)$ and $\chi_{q,t}(x_i) = Y_{i,q}(1 + V_{i,q^2})$.

Similarly we can compute x'_i . We have $\mathbb{L}(W) = L(W)$ always and the q -character is

$$\chi_{q,t}(x'_i)_{\leq 2} = \begin{cases} Y_{i,1}(1 + V_{i,q} \prod_j (1 + V_{j,q^2})^{a_{ij}}) & \text{if } i \in I_0, \\ Y_{i,q^3} & \text{if } i \in I_1. \end{cases}$$

This gives an answer to the exercise we mentioned after (5.2).

Proof of Theorem 6.3. Since $\mathbf{E}_W$ is a vector space by Proposition 4.6 and $IC_W(V)$'s are monodromic (i.e. $H^j(IC_W(V))$ is locally constant on every $\mathbb{C}^*$ -orbit of $\mathbf{E}_W$), we can apply the Fourier-Sato-Deligne functor Ψ ([36, 39]). For example, we have

$$\Psi(IC_W(0)) = 1_{\mathbf{E}_W^*}[\dim \mathbf{E}_W].$$

Other $\Psi(IC_W(V))$ are simple perverse sheaves on $\mathbf{E}_W^*$.

Recall that $\tilde{\mathcal{F}}(\nu, W)$ is a vector subbundle of the trivial bundle $\mathcal{F}(\nu, W) \times \mathbf{E}_W$ by Proposition 4.6. Let Ψ' be the Fourier-Sato-Deligne functor for this trivial bundle. We have

$$\Psi'(1_{\tilde{\mathcal{F}}(\nu, W)}[\dim \tilde{\mathcal{F}}(\nu, W)]) = 1_{\tilde{\mathcal{F}}(\nu, W)^\perp}[\dim \tilde{\mathcal{F}}(\nu, W)^\perp],$$

where $\tilde{\mathcal{F}}(\nu, W)^\perp$ is the annihilator in the dual trivial bundle $\mathcal{F}(V, W) \times \mathbf{E}_W^*$ as in §4.2. Moreover we have

$$\pi_1^\perp \circ \Psi' = \Psi \circ \pi_1.$$

Therefore if we decompose the pushforward as

$$\pi_!^\perp(1_{\tilde{\mathcal{F}}(\nu, W)^\perp}[\dim \tilde{\mathcal{F}}(\nu, W)^\perp]) \cong \bigoplus_{V', l} L_{V', l} \otimes \Psi(IC_W(V'))[l],$$

we have $\sum_l t^l \dim L_{V', l} = a_{V, V'; W}(t)$.

Take a general point of $\mathbf{E}_W^*$ and consider the Poincaré polynomial of the stalk of the above. In the left hand side we get the Poincaré polynomial of $\mathrm{Gr}_V(\sigma W)$ by Lemma 4.10. On the other hand, in the right hand side the factor $\Psi(IC_W(V'))$ with $IC_W(V') \notin \mathcal{L}_W$ disappears as its support is smaller than $\mathbf{E}_W^*$. For $IC_W(V') \in \mathcal{L}_W$, we get $r_W(IC_W(V')) \times a_{V, V'; W}(t)$, as $\Psi(IC_W(V'))$ is the IC complex associated with a local system of rank $r_W(IC_W(V'))$ defined over an open subset of $\mathbf{E}_W^*$. Thus we have

$$(6.6) \quad P_t(\mathrm{Gr}_V(\sigma W)) = \sum_{IC_W(V') \in \mathcal{L}_W} r_W(IC_W(V')) a_{V, V'; W}(t).$$

We get the assertion by recalling that $a_{V, V'; W}(t)$ is the coefficient of $e^{W^\perp} e^{V^\perp} = e^W e^V$ in the q -character of $L(W^\perp)$, where $\dim W^\perp = \dim W - \mathbf{C}_q \dim V'$, $\dim V^\perp = \dim V - \dim V'$ (§3.2). $\square$

6.3. Factorization of KR modules. In the remainder of this section, we give several simple applications of Theorem 6.3.

Proposition 6.7.

$$\mathbb{L}(W) \cong \mathbb{L}({}^\varphi W) \otimes \bigotimes_{i \in I} f_i^{\min(\dim W_i, \dim W_{i'})},$$

where ${}^\varphi W$ is given by

$$\dim {}^\varphi W_i = \max(\dim W_i - \dim W_{i'}, 0), \quad \dim {}^\varphi W_{i'} = \max(\dim W_{i'} - \dim W_i, 0)$$

The right hand side is independent of the order of the tensor product.

From this proposition it becomes enough to understand $\mathbb{L}({}^\varphi W)$. Notice that either ${}^\varphi W_i$ or ${}^\varphi W_{i'}$ is zero for each $i \in I$. If ${}^\varphi W_i = 0$, then ${}^\varphi W_{i'}$ is not connected to any other vertexes, and is easy to factor out it. Thus we eventually reduce to study the case when all ${}^\varphi W_{i'} = 0$, i.e. $\mathbf{E}_{{}^\varphi W}$ is the vector space of representations of the principal part of the decorated quiver obtained by deleting all frozen vertexes i' .

Proof. From the definition of σW in the formula (4.9) it is clear that σW is unchanged even if we add $\pm(1, 1)$ to $(\dim W_i, \dim W_{i'})$ for $i \in I_1$. Let $\tilde{W}$ be the $(I \sqcup I_{\mathrm{fr}})$ -graded vector space obtained from W by replacing both $W_i, W_{i'}$ by the vector space of dimension $\min(\dim W_i, \dim W_{i'})$ for each $i \in I_1$. Therefore we have

$$\chi_{q,t}(\mathbb{L}(W))_{\leq 2} = \chi_{q,t}(\mathbb{L}(\tilde{W}))_{\leq 2} \prod_{i \in I_1} (Y_{i,q} Y_{i,q^3})^{\min(\dim W_i, \dim W_{i'})}.$$

Since the truncated q -character of the Kirillov-Reshetikhin module is equal to $Y_{i,q} Y_{i,q^3}$ by Example 6.5, we have

$$\chi_{q,t}(\mathbb{L}(W))_{\leq 2} = \chi_{q,t}(\mathbb{L}(\tilde{W}))_{\leq 2} \prod_{i \in I_1} f_i^{\min(\dim W_i, \dim W_{i'})}.$$

Next we study a similar but slightly different reduction for $i \in I_0$. We consider the variety $(\pi^\perp)^{-1}(\bigoplus \mathbf{x}_i^*, \bigoplus \mathbf{y}_h^*)$ as in the statement of Lemma 4.10. $((\bigoplus \mathbf{x}_i^*, \bigoplus \mathbf{y}_h^*))$ is a representation before applying the reflection functors as in the proof of Lemma 4.10.) From the condition $X_i \subset$

$\text{Ker } x_i^*$ it is isomorphic to $(\bar{\pi}^\perp)^{-1}(\bigoplus \bar{\mathbf{x}}_i^*, \bigoplus \bar{\mathbf{y}}_h^*)$, where (1) $\tilde{W}$ obtained from W by replacing W_i by $\text{Ker } x_i^*$, (2) $\bar{\mathbf{y}}_h^*$ is the restriction of $\mathbf{y}_h^*$ and other maps are obvious ones. We have

$$\dim \tilde{W}_i = \max(\dim W_i - \dim W_{i'}, 0).$$

Therefore we have

$$\chi_{q,t}(\mathbb{L}(W))_{\leq 2} = \chi_{q,t}(\mathbb{L}(\tilde{W}))_{\leq 2} \prod_{i \in I_0} (Y_{i,1} Y_{i,q^2})^{\min(\dim W_i, \dim W_{i'})}.$$

Note again that $Y_{i,1} Y_{i,q^2}$ is the truncated q -character of the Kirillov-Reshetikhin module f_i . Therefore the above equality can be written as

$$\chi_{q,t}(\mathbb{L}(W))_{\leq 2} = \chi_{q,t}(\mathbb{L}(\tilde{W}))_{\leq 2} \prod_{i \in I_0} f_i^{\min(\dim W_i, \dim W_{i'})}.$$

Combining these two reductions we obtain the assertion. $\square$

6.4. Factorization and canonical decomposition. Take a general representation $(\bigoplus \mathbf{y}_h)$ of $\mathbf{E}_{\varphi W}$. We decompose it into a sum of indecomposable representations. We have a corresponding decomposition

$$\varphi W = W^1 \oplus W^2 \oplus \dots \oplus W^s$$

of the I -graded graded vector space. It is known [34, p.85] that $W^1, \dots, W^s$ are independent of a choice of general representation of $\mathbf{E}_{\varphi W}$ up to permutation. This is called the *canonical decomposition* of φW (or $\dim \varphi W$). It is known that all $\dim W^\alpha \in \mathbb{Z}_{\geq 0}^I$ are *Schur roots* and $\text{ext}^1(W^k, W^l) = 0$ for $k \neq l$ ([35, Prop. 3]). Here $\dim W^k$ is a Schur root if a general representation in $\mathbf{E}_{W^k}$ has only trivial endomorphisms, i.e. scalars. It is known that this is equivalent to a general representation is indecomposable ([loc. cit., Prop. 1]). And $\text{ext}^1(W^k, W^l)$ is the dimension of Ext^1 between general representations in $\mathbf{E}_{W^k}$ and $\mathbf{E}_{W^l}$. Basic results on the canonical decomposition were obtained by Schofield [56], which will be used in part below.

Note that the frozen part play no role in the canonical decomposition, as $\varphi W_{i'} \neq 0$ implies $\varphi W_i = 0$. Therefore we simply have factors $\underbrace{S_{i'} \oplus \dots \oplus S_{i'}}_{\dim \varphi W_{i'} \text{ factors}}$ in the canonical decomposition. If

φW contains a factor $S_i^{\oplus m_i}$ for $i \in I_1$, it is killed by $\sigma(\cdot)$. We thus have

Proposition 6.8. *Suppose that the canonical decomposition of φW contains factors as*

$$\varphi W = \psi W \oplus \bigoplus_{i \in I} S_{i'}^{\oplus \dim \varphi W_{i'}} \oplus \bigoplus_{i \in I_1} S_i^{\oplus m_i}.$$

Then we have a factorization

$$\mathbb{L}(\varphi W) = \mathbb{L}(\psi W) \otimes \bigotimes_{i \in I} L(S_{i'})^{\otimes \dim \varphi W_{i'}} \otimes \bigotimes_{i \in I_1} L(S_i)^{\otimes m_i}.$$

We consider the following condition (C):

(C) The canonical decomposition of φW contains only *real* Schur roots.

Proposition 6.9. (1) *Assume the condition (C). Then $\mathcal{L}_{\varphi W} = \{IC_{\varphi W}(0)\}$ and hence $\mathbb{L}(W) = L(W)$.*

(2) *If $\mathcal{L}_{\varphi W} = \{IC_{\varphi W}(0)\}$, $\text{Gr}_V(\sigma W)$ has no odd cohomology group.*

Proof. (1) From the definition, $\mathbf{E}_{\varphi W}$ contains the $\prod_i \mathrm{GL}(W_i) \times \mathrm{GL}(W_{i'})$ orbit of a general representation as a Zariski open subset. The same is true for $\mathbf{E}_{\varphi W}^*$. Since all $\Psi(IC_W(V))$ are $\prod_i \mathrm{GL}(W_i) \times \mathrm{GL}(W_{i'})$ -equivariant, we cannot have IC complexes associated with nontrivial local systems as stabilizers are always connected. Therefore we only have $\mathcal{L}_{\varphi W} = \{IC_{\varphi W}(0)\}$.

(2) Since (6.6) is a single sum, the assertion follows from Remark 3.20. $\square$

Proposition 6.10.

$$(6.11) \quad \mathbb{L}({}^\varphi W) \cong \mathbb{L}(W^1) \otimes \cdots \otimes \mathbb{L}(W^s).$$

Proof. We assume $s = 2$. Since we do not use the assumption that W^1, W^2 are Schur roots, the proof also gives the proof for general case.

Consider the convolution diagram in §3.5. By [43, 10.1] the restriction functor commutes with the Fourier-Sato-Deligne functor up to shift. Therefore we consider perverse sheaves defined over $\mathbf{E}_{W^1}^*, \mathbf{E}_{W^2}^*, \mathbf{E}_W^*$.

We take open subsets U^1, U^2 in $\mathbf{E}_{W^1}^*, \mathbf{E}_{W^2}^*$ so that perverse sheaves not in $\mathcal{L}_{W^1}, \mathcal{L}_{W^2}$ have support outside of U^1, U^2 . Similarly we take an open subset $U \subset \mathbf{E}_W^*$ consisting of modules isomorphic to direct sum of modules in U^1 and U^2 , and perverse sheaves not in $\mathcal{L}_W$ have support outside of U .

We may assume that Ext-groups between modules in U^1, U^2 vanish. Therefore any module in $\kappa^{-1}(U^1 \times U^2)$ is isomorphic to direct sum of modules from U^1 and U^2 . Therefore $\kappa^{-1}(U^1 \times U^2) \subset U$ and κ is an isomorphism. Therefore for $L \in \mathcal{P}_W \setminus \mathcal{L}_W$, $\mathrm{Res} L$ does not have factors in $IC_{W^1}(V^1) \boxtimes IC_{W^2}(V^2)$ with $IC_{W^\alpha}(V^\alpha) \in \mathcal{L}_{W^\alpha}$ ($\alpha = 1, 2$). Therefore the product of $L(W^1) \in \mathcal{L}_{W^1}$ and $L(W^2) \in \mathcal{L}_{W^2}$ is a linear combination of elements in $\mathcal{L}_W$.

If $IC_W(V) \in \mathcal{L}_W$, the restriction of $\kappa_! \iota^* \Psi(IC_W(V))$ to $U^1 \times U^2$ is a local system of rank $r(IC_W(V))$. Thus if we write

$$\begin{aligned} \mathrm{Res} IC_W(V) = & \sum_{IC_{W^1}(V^1) \in \mathcal{L}_{W^1}, IC_{W^2}(V^2) \in \mathcal{L}_{W^2}} a_V^{V^1, V^2} IC_{W^1}(V^1) \boxtimes IC_{W^2}(V^2) \\ & + (\text{linear combination of } L \in \mathcal{P}_W \setminus \mathcal{L}_W), \end{aligned}$$

then $a_V^{V^1, V^2}$ is an integer (up to shift). And we have

$$r(IC_W(V)) = \sum_{V^1, V^2} a_V^{V^1, V^2} r(IC_{W^1}(V^1)) r(IC_{W^2}(V^2)).$$

From this we have $\mathbb{L}(W^1) \otimes \mathbb{L}(W^2) = \mathbb{L}(W)$. $\square$

Let us show the converse.

Proposition 6.12. (1) Suppose that $\mathbb{L}({}^\varphi W)$ decomposes as

$$\mathbb{L}({}^\varphi W) \cong \mathbb{L}(W^1) \otimes \mathbb{L}(W^2).$$

Then we have $\mathrm{ext}^1(W^1, W^2) = 0 = \mathrm{ext}^1(W^2, W^1)$.

(2) The factorization of an almost simple module $\mathbb{L}(W)$ is exactly given by the canonical decomposition of ${}^\varphi W$, and we have the bijection

$$\left\{ \begin{array}{l} \text{prime almost simple modules} \\ \text{with (C)} \end{array} \right\} \setminus \{x_i, f_i \mid i \in I\} \longleftrightarrow \left\{ \begin{array}{l} \text{Schur roots of the principal} \\ \text{part } \mathcal{Q} \text{ of the decorated quiver} \end{array} \right\}$$

given by $\mathbb{L}(W) \leftrightarrow \dim W$.

Here an almost prime simple module $\mathbb{L}(W)$ means that it does not factor as $\mathbb{L}(W^1) \otimes \mathbb{L}(W^2)$ of almost simple modules.

Proof. Let us first consider the case $x_{i'} = L(W^2)$. Taking the truncated q -character, we have

$$\sum_V P_t(\mathrm{Gr}_V(\sigma W)) e^W e^V = Y_{i,q^3} * \left(\sum_{V^1} P_t(\mathrm{Gr}_{V^1}(\sigma W^1)) e^{W^1} e^{V^1} \right),$$

where $*$ is the twisted multiplication (3.21).

Since $i \in I_1$ is a source, we have $\mathrm{ext}^1(W^1, S_i) = 0$. If we have $\mathrm{ext}^1(S_i, W^1) \neq 0$, then $\dim \sigma W^1 = \dim \sigma W + \dim S_i$. Therefore the right hand side contains the term for V^1 with $\dim V^1 = \dim \sigma W + \dim S_i$, as the corresponding quiver Grassmannian $\mathrm{Gr}_{\sigma W^1}(\sigma W^1)$ is a single point.

But the left hand side obviously cannot contain the corresponding term. Therefore we must have $\mathrm{ext}^1(S_i, W^1) = 0$.

Now we suppose generic representations of W^1 and W^2 do not contain the direct summand S_i for any $i \in I_1$. Then the vanishing of ext^1 is equivalent to the corresponding statement after applying the functor σ . (Since σ starts with taking the *dual*, we need to exchange the first and the second entries A, B of $\mathrm{ext}^1(A, B)$, but we are studying both $\mathrm{ext}^1(A, B)$ and $\mathrm{ext}^1(B, A)$, so it does not matter.)

We again consider the equality for the truncated q -character:

$$\sum_V P_t(\mathrm{Gr}_V(\sigma W)) e^W e^V = \left(\sum_{V^1} P_t(\mathrm{Gr}_{V^1}(\sigma W^1)) e^{W^1} e^{V^1} \right) * \left(\sum_{V^2} P_t(\mathrm{Gr}_{V^2}(\sigma W^2)) e^{W^2} e^{V^2} \right).$$

The right hand side contain the terms with $V^1 = \sigma W^1$, $V^2 = 0$ and $V^1 = 0$, $V^2 = \sigma W^2$, as both $\mathrm{Gr}_{V^1}(\sigma W^1)$ and $\mathrm{Gr}_{V^2}(\sigma W^2)$ are points in these cases. These survive thanks to the positivity $P_t(\mathrm{Gr}_{V^1}(\sigma W^1)), P_t(\mathrm{Gr}_{V^2}(\sigma W^2)) \in \mathbb{Z}_{\geq 0}[t]$. Therefore the corresponding quiver Grassmannian varieties $\mathrm{Gr}_V(\sigma W)$ (two cases) are nonempty in the left hand side also. Therefore a general representation of $\mathbf{E}_W$ contains two subrepresentations of $\dim \sigma W^1$, $\dim \sigma W^2$ respectively. By [56, Th. 3.3], it implies that we have both $\mathrm{ext}^1(\sigma W^1, \sigma W^2) = 0$ and $\mathrm{ext}^1(\sigma W^2, \sigma W^1) = 0$. This proves the first assertion.

The second assertion follows from the first and the characterization of the canonical decomposition: $\alpha = \sum \beta^i$ is the canonical decomposition if and only if each β^i is a Schur root and $\mathrm{ext}^1(\beta^i, \beta^j) = 0$ for $i \neq j$. (See [35, Prop. 3].) $\square$

Corollary 6.13. *If $L(W)$ satisfies (C), it is real, i.e. $L(W) \otimes L(W)$ is simple.*

At this moment, we do not know the converse is true or not.

Next suppose $\mathcal{G}$ is of type *ADE*. Then all positive roots are real and Schur. Let Δ_+ be the set of positive roots. Following [22] we introduce the set $\Phi_{\geq -1}$ of *almost positive* roots:

$$\Phi_{\geq -1} = \Delta_+ \sqcup \{-\alpha_i \mid i \in I\},$$

where α_i is the simple root for i .

Corollary 6.14. (1) *There are only finitely many prime simple modules in $\mathbf{R}_{\ell=1}$ if and only if the underlying graph $\mathcal{G}$ of the principal part is of type *ADE*.*

(2) *Suppose that $\mathcal{G}$ is of type *ADE*. Then all simple modules are real, and there is a bijection*

$$\{\text{prime simple modules}\} \setminus \{f_i \mid i \in I\} \xrightarrow[1:1]{\dim(\bullet)} \Phi_{\geq -1}.$$

Here the bijection is given by Proposition 6.12(2) together with $x_i \mapsto (-\alpha_i)$.

The first assertion is a simple consequence of the fact that there are infinitely many real Schur roots for non *ADE* quivers. This can be shown for example, by observing non *ADE* graph always contains an affine graph. Then for an affine graph, real roots α with the defects $\chi(\delta, \alpha) = \dim \operatorname{Hom}(\delta, \alpha) - \dim \operatorname{Ext}^1(\delta, \alpha)$ are nonzero are Schur. Here δ is the generator of positive imaginary roots and the above is Euler form for a representation N with $\dim N = \delta$ and M with $\dim M = \alpha$, which is independent of the choice of M, N . This result is nothing but [22] after identifying prime simple modules with cluster variables in the next section.

Now we consider the affine case.

Example 6.15. Suppose that (I, E) of type $A_1^{(1)}$. The corresponding quiver (I, Ω) is called the Kronecker quiver. Positive roots are $(n \rightrightarrows n+1), (n+1 \rightrightarrows n), (n \rightrightarrows n) (n \in \mathbb{Z}_{\geq 0})$. The vector $(1 \rightrightarrows 1)$ is the generator of positive imaginary roots, and denoted by δ as usual.

For $n \in \mathbb{Z}_{>0}$ let nW denote an $(I \sqcup I_{\text{fr}})$ -graded vector space with $\mathbb{C}^n$ at the entry i and 0 at i' ($i = 0, 1$): $(nW)_0 = \mathbb{C}^n \rightrightarrows (nW)_1 = \mathbb{C}^n$. Thus $\dim(nW) = n\delta$. Then $nW = W \oplus \cdots \oplus W$ is the canonical decomposition of nW , where W means $1W$: It is well-known that a general representation in $\mathbf{E}_W$ corresponds to a point in $\mathbb{P}^1(\mathbb{C})$. And a general representation in $\mathbf{E}_{nW}$ corresponds to distinct n points in $\mathbb{P}^1(\mathbb{C})$.

For a real positive root $(n \rightrightarrows n+1)$ or $(n+1 \rightrightarrows n)$, there is the unique indecomposable module M . It is known that either $\operatorname{Ext}^1(M, W)$ or $\operatorname{Ext}^1(W, M)$ are nonvanishing. Therefore M and W cannot appear in a canonical decomposition simultaneously. It is also known that extensions between $(n \rightrightarrows n+1)$ and $((n+1) \rightrightarrows (n+2))$ vanish. It is also true for $((n+1) \rightrightarrows n)$ and $((n+2) \rightrightarrows (n+1))$. All other pairs, one of extensions does not vanish.

From these observations, the canonical decomposition only have real Schur roots, except the case nW . We consider the case $n = 2$. If we consider $\pi^\perp: \tilde{\mathcal{F}}(\nu, 2W)^\perp \rightarrow \mathbf{E}_W^*$ in §4.2, the perverse sheaves appearing (up to shift) in the pushforward $\pi_!^\perp(1_{\tilde{\mathcal{F}}(\nu, 2W)^\perp}[\dim \tilde{\mathcal{F}}(\nu, 2W)^\perp])$ was studied in [42]. If we take $\nu = (1, 1) \in \mathbb{Z}_{\geq 0}^I$, then $\pi^\perp$ is the principal $\{\pm 1\}$ cover over the open set $\mathbf{E}_W^{*\text{reg}}$ corresponding to distinct pairs of points in $\mathbb{P}^1(\mathbb{C})$. Then from [loc. cit.] we have

$$\mathcal{L}_{2W} = \{1_{\{0\}}, \Psi^{-1}(IC(\mathbf{E}_{2W}^*, \rho))\},$$

where $IC(\mathbf{E}_{2W}^*, \rho)$ is the IC complex associated with the nontrivial local system ρ corresponding to the nontrivial representation of $\{\pm 1\}$. In particular, the almost simple module $\mathbb{L}(2W)$ is not the simple module $L(2W)$. On the other hand $\mathcal{L}_W = \{1_{\{0\}}\}$.

The coefficient of $\chi_q(L(2W))$ at $Y_{1,1}^2 Y_{2,q^3}^2 \times V_{1,q} V_{2,q^2}$ is 1. The coefficients of $\chi_q(L(W))$ at $Y_{1,1} Y_{2,q^3} V_{1,q}, Y_{1,1} Y_{2,q^3} V_{2,q^2}$ are both 1. Therefore $L(2W) \not\cong L(W) \otimes L(W)$, i.e. $L(W)$ is *not* real. On the other hand, we have $\mathbb{L}(2W) \cong \mathbb{L}(W) \otimes \mathbb{L}(W)$.

There are many attempts to construct a base for the cluster algebra corresponding to this example in the cluster algebra literature ([57, 10, 20, 19] and [26] in a wider context). The problem is how to understand imaginary root vectors, and the solution is not unique. Relationship between various bases are studied by Leclerc [41].

More generally if W corresponds to an indivisible isotropic imaginary root (i.e. in the Weyl group orbit of δ of a subdiagram of affine type in $\mathcal{G}$) in an arbitrary $\mathcal{Q}$, we have

$$\mathbb{L}(nW) \cong \mathbb{L}(W)^{\otimes n}.$$

This can be generalized thanks to [56]. First we have if α is a non-isotropic imaginary Schur root, $n\alpha$ is also a Schur root for $n \in \mathbb{Z}_{>0}$ ([loc. cit., Th. 3.7]). It is also known that an isotropic Schur root must be indivisible ([loc. cit., Th. 3.8].) Therefore we introduce the following notation: For a W as above and $n \in \mathbb{Z}_{>0}$ let nW be an I -graded vector space with

$\dim(nW)_i = n \dim W_i$. For a factor $\mathbb{L}(W^k)$ in (6.11) let $(n\mathbb{L})(W^k)$ be $\mathbb{L}(nW^k)$ if $\dim W^k$ is a non-isotropic Schur imaginary root, and $\mathbb{L}(W^k)^{\otimes n}$ otherwise, i.e. $\dim W^k$ is a real or indivisible isotropic Schur root.

Corollary 6.16. *Let W be as above. Let $\varphi W = W^1 \oplus W^2 \oplus \cdots \oplus W^s$ be the canonical decomposition. Then we have*

$$\mathbb{L}(nW) \cong (n\mathbb{L})(W^1) \otimes \cdots \otimes (n\mathbb{L})(W^s) \otimes \bigotimes_{i \in I} f_i^{n \min(\dim W_i, \dim W_{i'})}.$$

Since $\mathbb{L}(2W) \not\cong \mathbb{L}(W) \otimes \mathbb{L}(W)$ for W corresponding non-isotropic imaginary Schur root, it is natural to hope the same is true for $L(2W)$ and $L(W) \otimes L(W)$.

7. CLUSTER ALGEBRA STRUCTURE

In this section we prove that cluster monomials are dual canonical base elements after some preparation.

In the previous sections, we use the notation W for an $(I \sqcup I_{\text{fr}})$ -graded representation. In this section we also use it for its general representation. Or if we first take a representation, its underlying $(I \sqcup I_{\text{fr}})$ -graded vector space will be denoted by the same notation.

7.1. Tilting modules. We first review the theory of tilting modules. (See [1, VI] and [28].)

Let $\mathcal{Q} = (I, \Omega)$ be a quiver as in §2. Let $\mathbb{C}\mathcal{Q}$ be its path algebra defined over $\mathbb{C}$. We consider the category $\text{rep}\mathcal{Q}$ of finite dimensional representations of $\mathcal{Q}$ over $\mathbb{C}$, which is identified with the category of finite dimensional $\mathbb{C}\mathcal{Q}$ -modules.

A module M of the quiver is said to be a *tilting module* if the following two conditions are satisfied:

- (1) M is *rigid*, i.e. $\text{Ext}^1(M, M) = 0$.
- (2) There is an exact sequence $0 \rightarrow \mathbb{C}\mathcal{Q} \rightarrow M_0 \rightarrow M_1 \rightarrow 0$ with $M_0, M_1 \in \text{add}M$, where $\text{add}M$ denotes the additive category generated by the direct summands of M .

We usually assume M is multiplicity free.

It is known that the number of indecomposable summands of M equals to the number of vertexes $\#I$, i.e. $\text{rank of } K_0(\mathbb{C}\mathcal{Q})$.

An rigid module M always has a module X so that $M \oplus X$ is a tilting module.

A module M is said to be an *almost complete tilting module* if it is rigid and the number of indecomposable summands of M is $\#I - 1$. We say an indecomposable module X is *complement* of M if $M \oplus X$ is a tilting module.

We have the following structure theorem:

Theorem 7.1 (Happel-Unger [28]). *Let M be an almost complete tilting module.*

(1) *If M is sincere, there exists two nonisomorphic complements X, Y which are related by an exact sequence*

$$0 \rightarrow X \rightarrow E \rightarrow Y \rightarrow 0$$

with $E \in \text{add}M$. Moreover, we have $\text{Ext}^1(Y, X) \cong \mathbb{C}$, $\text{Ext}^1(X, Y) = 0$, $\text{Hom}(Y, X) = 0$.

(2) *If M is not sincere, there exists only one complement X up to isomorphisms.*

Here a module M is said to be *sincere* if $M_i \neq 0$ for any vertex i .

7.2. Cluster tilting sets. When the quiver $\mathcal{Q}$ does not contain an oriented cycle (i.e. acyclic quiver), combinatorics of the cluster algebra can be understood from the cluster category theory. Since we only need the statement, we explain the theory only very briefly following [32]. We only consider the case when there are no frozen variables.

Let $n = \#I$. A collection $\mathbf{L} = \{W^1, \dots, W^n\}$ is said to be a *cluster-tilting set* if the following conditions are satisfied:

- (0) W^i is either an indecomposable representation of the quiver $\mathcal{Q}$ or a vertex. Let $\mathbf{L}_{\text{mod}}$ be the subset of indecomposable representations, $\mathbf{L}_{\text{ver}} = \mathbf{L} \setminus \mathbf{L}_{\text{mod}}$.
- (1) $W^k \in \mathbf{L}_{\text{mod}}$ are pairwise nonisomorphic. $W^i \in \mathbf{L}_{\text{ver}}$ are pairwise distinct.
- (2) Delete all arrows incident with a vertex $W^i \in \mathbf{L}_{\text{ver}}$. Remove the vertex W^i . Let ${}^\psi\mathcal{Q}$ be the resulted quiver.
- (3) The entry for $W^k \in \mathbf{L}_{\text{mod}}$ is 0 for a vertex $W^j \in \mathbf{L}_{\text{ver}}$. Hence W^k is a representation of ${}^\psi\mathcal{Q}$.
- (4) ${}^\psi W \stackrel{\text{def.}}{=} \bigoplus_{W^k \in \mathbf{L}_{\text{mod}}} W^k$ is a tilting module as a representation of ${}^\psi\mathcal{Q}$.

Note that $\#\mathbf{L}_{\text{mod}} = \#({}^\psi I)$. Therefore ${}^\psi W$ is tilting if and only if $\text{ext}^1(W^k, W^l) = 0$ for any k, l (including the case $k = l$). Thus this is stronger than the canonical decomposition and means that $\dim W^k$ is a real Schur root.

The initial cluster-tilting set is the collection $\mathbf{L} = I$ with $\mathbf{L}_{\text{mod}} = \emptyset$. In this case ${}^\psi I = \emptyset$ and the condition is trivially satisfied.

If we identify $W^i \in \mathbf{L}_{\text{ver}}$ with $P_{W^i}[1]$ the shift of the indecomposable projective module associated corresponding to the vertex W^i , the above definition is nothing but the definition of a cluster-tilting set for the cluster category [6].

For $k \in \{1, \dots, n\}$ we define the *mutation* $\mu_k(\mathbf{L})$ of $\mathbf{L}$ in direction k as follows:

- (1) Suppose W^k is a vertex. We return back it and all arrows incident with it to the quiver ${}^\psi\mathcal{Q}$. Let ${}^{+\psi}\mathcal{Q}$ be the resulted quiver. Since ${}^\psi W$ is an almost tilting non-sincere module as a representation of ${}^{+\psi}\mathcal{Q}$, we can add the unique indecomposable $*W^k$ to ${}^\psi W$ to get a tilting module.
- (2) Next suppose W^k is a module. We consider an almost tilting module ${}^{-\psi}W$ which is obtained from ${}^\psi W$ by subtracting the summand W^k .
 - (a) If it is sincere, there is another indecomposable module $*W^k \neq W^k$ such that $*W^k \oplus {}^{-\psi}W$ is a tilting module.
 - (b) If it is not sincere, there exists the unique simple module S_i , not appearing in the composition factors of ${}^{-\psi}W$. Then we set $*W^k = i$.

Let

$$\mu_k(\mathbf{L}) \stackrel{\text{def.}}{=} \mathbf{L} \cup \{*W^k\} \setminus \{W^k\}.$$

In all cases $\mu_k(\mathbf{L})$ is again a cluster-tilting set. We can iterate this procedure and obtain new clusters starting from the initial cluster $\mathbf{L} = I$.

7.3. Cluster character. We still continue to assume that the quiver $\mathcal{Q}$ does not contain an oriented cycle. It is known that cluster monomials can be expressed in terms of generating functions of Euler numbers of quiver Grassmannian varieties. This important result was first proved by Caldero-Chapoton in type *ADE* [7]. Later it was generalized to any acyclic quiver by Caldero-Keller [8] using various results in the cluster category theory (see [37] for the reference). We recall the formula in this subsection.

Let $(\mathbf{x}, \mathbf{B})$ be the initial seed of the cluster algebra $\mathcal{A}(\mathbf{B})$. We assume there is no frozen part for simplicity. Let W be a representation of the quiver $\mathcal{Q}$ corresponding to $\mathbf{B}$. Let $\text{Gr}_V(W)$ be

the corresponding quiver Grassmannian variety, where V is an I -graded vector space. Though we soon assume W is a general representation in $\mathbf{E}_W$, it is not necessary for the definition. Let $e(\mathrm{Gr}_V(W))$ be its Euler number. We define

$$X_W \stackrel{\text{def.}}{=} \frac{1}{\mathbf{x}^{\dim W}} \sum_V e(\mathrm{Gr}_V(W)) \mathbf{x}^{\dim V \cdot R} \mathbf{x}^{(\dim W - \dim V)R'},$$

where

$$\begin{aligned} \mathbf{x}^{\dim W} &= \prod_i x_i^{\dim W_i}, \\ \mathbf{x}^{\dim V \cdot R} &= \prod_{h \in \Omega} x_{i(h)}^{\dim V_{o(h)}}, \quad \mathbf{x}^{(\dim W - \dim V)R'} = \prod_{h \in \Omega} x_{o(h)}^{(\dim W_{i(h)} - \dim V_{i(h)})}. \end{aligned}$$

For a vertex i , we set $X_i = x_i$.

Then it is known that the correspondence $W \rightarrow X_W$ gives the followings:

- the correspondence $W \rightarrow X_W$ defines a bijection between the set of isomorphism classes of rigid indecomposable modules with cluster variables minus $\{x_i\}$;
- the correspondence $\mathbb{L} \rightarrow \{X_{W^1}, \dots, X_{W^n}\}$ gives a bijection between cluster tilting sets and clusters;
- the mutation on cluster tilting sets corresponds to the cluster mutation.

7.4. Piecewise-linear involution. We give one more preparation before applying results from the cluster category theory to our setting. This last preliminary is not necessary for our argument, but helps to make a relation to [31, §12.3].

We recall the piecewise-linear involution τ_- on the root lattice considered in [31, §7]: for $\gamma = \sum_i \gamma_i i \in \mathbb{Z}^I$, we define $\tau_-(\gamma) = \sum_i \tau_-(\gamma)_i i$ by

$$(7.2) \quad \tau_-(\gamma)_i = \begin{cases} -\gamma_i - \sum_{j \neq i} c_{ij} \max(0, \gamma_j) & \text{if } i \in I_1, \\ \gamma_i & \text{if } i \in I_0, \end{cases}$$

where (c_{ij}) is the Cartan matrix.

Let

$$(7.3) \quad \gamma = \sum_i (\dim W_i - \dim W_{i'}) i.$$

If $i \in I_0$, we have

$$\tau_-(\gamma)_i = \dim W_i - \dim W_{i'} = \dim {}^\varphi W_i - \dim {}^\varphi W_{i'}.$$

If $i \in I_1$, we have

$$\begin{aligned} \tau_-(\gamma)_i &= \dim W_{i'} - \dim W_i - \sum_{j \neq i} c_{ij} \max(\dim W_j - \dim W_{j'}, 0) \\ &= \dim {}^\varphi W_{i'} - \dim {}^\varphi W_i - \sum_{j \neq i} c_{ij} \dim {}^\varphi W_j. \end{aligned}$$

Therefore we have

$$\dim {}^{\sigma\varphi} W_i = \max(\tau_-(\gamma)_i, 0).$$

where ${}^{\sigma\varphi} W = \sigma({}^\varphi W)$ is obtained by applying σ to ${}^\varphi W$.

Remark 7.4. In [31, §12.3] the quiver Grassmannian $\text{Gr}_V(M[\tau_-(\gamma)])$ was considered where $M[\tau_-(\gamma)]$ is a generic representation with

$$\dim M[\tau_-(\gamma)]_i = \max(\tau_-(\gamma)_i, 0).$$

Here the quiver is the principal part $\mathcal{Q}$ of our decorated quiver. From the above computation, we have $M[\tau_-(\gamma)]$ is nothing but the principal quiver part of ${}^\sigma W$. The frozen part of ${}^\sigma W$ does not play any role in the quiver Grassmannian, by Proposition 6.8. Therefore $\text{Gr}_V(M[\tau_-(\gamma)])$ in [loc. cit., §12.3] and our $\text{Gr}_V({}^\sigma W)$ is isomorphic under (7.3).

7.5. Cluster monomials. We start to put the cluster algebra structure on $\mathbf{R}$ from this subsection.

Proposition 7.5. (1) *Let W be an I -graded vector space such that $\dim W$ is a real Schur root of the principal part of the decorated quiver. Then $L(W)$ is a cluster variable.*

(2) *This correspondence defines a bijection between the set of real Schur roots and the set of cluster variables except variables in the initial seed, i.e. x_i, f_i ($i \in I$).*

For type ADE , this together with Corollary 6.14 shows the condition (2) in the monoidal categorification 2.4.

Proof. Roughly this is a consequence of results reviewed in §7.3. However, our quiver Grassmannian is for ${}^\sigma W$, not for W . Correspondingly we need to replace the initial seed of $\mathcal{A}(\tilde{\mathbf{B}})$ by the $\mathbf{z}$ -quiver in (5.8). When we mutate from $\mathbf{x}$ -quiver to $\mathbf{z}$ -quiver, the set of cluster variables does not change by definition, but variables in the initial seed change. So let us first consider this effect. The functor ${}^\sigma(\bullet)$ induces an involution on the set

$$\{\text{real Schur roots}\} \setminus \{\alpha_i \mid i \in I_1\}.$$

Therefore we only need to study cluster variables corresponding to α_i in either $\mathbf{x}$ -quiver or $\mathbf{z}$ -quiver.

- In $\mathbf{x}$ -quiver, α_i corresponds to $W = S_i$. We have $L(S_i) = x'_i = z_i$. This is a cluster variable of the seed for the $\mathbf{z}$ -quiver, but not for the original $\mathbf{x}$ -quiver. Note also that ${}^\sigma W = 0$ in this case.
- In $\mathbf{z}$ -quiver, α_i corresponds to the cluster variable obtained as z_i^* . But this is nothing but x_i . The corresponding simple module is $L(S_{i'})$. We do not consider since it has support in the frozen part.

We now may assume $\dim {}^\sigma W$ is a real Schur root different from α_i ($i \in I_1$).

We cannot apply the formula in §7.3 directly as the $\mathbf{z}$ -quiver contains an oriented cycle in general. (See (5.7).) We thus first consider the quiver with principal coefficients, and write down F -polynomials and $\mathbf{g}$ -vectors by using the formula in §7.3. Then we apply the result in §2.2 to get the formula for cluster variables in the original cluster algebra.

We take $\mathbf{u}, \mathbf{f}$ as cluster variables for the initial seed of $\mathcal{A}_{\text{pr}}$ and define

$$X_{\sigma W}(\mathbf{u}, \mathbf{f}) \stackrel{\text{def.}}{=} \frac{1}{\prod_{i \in I} u_i^{\sigma w_i}} \sum_V e(\text{Gr}_V({}^\sigma W)) \prod_{i \in I_0} u_i^{\sum_j a_{ij} v_j} \prod_{i \in I_1} u_i^{\sum_j a_{ij} ({}^\sigma w_j - v_j)} \prod_{i \in I} f_i^{v_i},$$

where $v_i = \dim V_i$, $w_i = \dim W_i$, $\sigma w_i = \dim {}^\sigma W_i$. By §7.3 this is a cluster variable α for $\mathcal{A}_{\text{pr}}$, and hence above gives the Laurent polynomial $X_\alpha(\mathbf{u}, \mathbf{f})$ in §2.2.

Hence the F -polynomial is

$$F_{\sigma W}(\mathbf{f}) = \sum_V e(\text{Gr}_V({}^\sigma W)) \prod_{i \in I} f_i^{v_i}.$$

And the $\mathbf{g}$ -vector is

$$\mathbf{g}_{\sigma W} = - \sum_{i \in I_0} \sigma w_i i - \sum_{i \in I_1} \left(\sigma w_i - \sum_j a_{ij} \sigma w_j \right) i = - \sum_i w_i \varepsilon_i i.$$

Now we return back to our original cluster algebra. Since our initial seed is given by the $\mathbf{z}$ -quiver, we change the notation in §2.2 and use z -variables instead of x -variables. We denote the cluster variable corresponding to above $X_{\sigma W}$ by $z[\sigma W]$. We have

$$z[\sigma W] = \frac{F_{\sigma W}(\widehat{\mathbf{y}})}{F_{\sigma W}|_{\mathbb{P}}(\mathbf{y})} \mathbf{z}^{\mathbf{g}_{\sigma W}},$$

where

$$y_j = \begin{cases} f_j^{-1} \prod_{i \in I} f_i^{a_{ij}} & \text{if } j \in I_0, \\ f_j^{-1} & \text{if } j \in I_1, \end{cases} \quad \widehat{y}_j = y_j \prod_{i \in I} z_i^{\varepsilon_i a_{ij}} \quad (j \in I).$$

in this situation. A direct calculation shows (see [31, Lem. 7.2])

$$\chi_q(\widehat{y}_j) = V_{j, q^{\varepsilon_j + 1}}.$$

We note that $F_{\sigma W}$ contains the monomial $\prod_i f_i^{\sigma w_i}$ for $V = \sigma W$ with the coefficient 1, and all other terms are its factor. If we evaluate it at y_j , we have

$$\prod_{i \in I} f_i^{-\sigma w_i} \prod_{i \in I_1} f_i^{\sum_j a_{ij} \sigma w_j} = \prod_{i \in I_0} f_i^{-\sigma w_i} \prod_{i \in I_1} f_i^{-\sigma w_i + \sum_j a_{ij} \sigma w_j} = \prod_{i \in I_0} f_i^{-w_i} \prod_{i \in I_1} f_i^{w_i}.$$

We also have the constant term 1 for $V = 0$. Therefore

$$F_{\sigma W}|_{\mathbb{P}}(\mathbf{y}) = \prod_{i \in I_0} f_i^{-w_i}.$$

Thus combining with the above calculation of $\mathbf{g}_{\sigma W}$, we get ([31, Lem. 7.3])

$$\frac{\mathbf{z}^{\mathbf{g}_{\sigma W}}}{F_{\sigma W}|_{\mathbb{P}}(\mathbf{y})} = \prod_{i \in I_0} f_i^{w_i} \prod_{i \in I} z_i^{-w_i \varepsilon_i}.$$

Its q -character is

$$\chi_q \left(\frac{\mathbf{z}^{\mathbf{g}_{\sigma W}}}{F_{\sigma W}|_{\mathbb{P}}(\mathbf{y})} \right) = \prod_{i \in I_0} Y_{i,1}^{w_i} \prod_{i \in I_1} Y_{i,q^3}^{w_i}.$$

We thus get

$$\chi_q(z[\sigma W]) = \sum_V e(\mathrm{Gr}_V(\sigma W)) e^W e^V.$$

Hence we have $z[\sigma W] = \mathbb{L}(W) = L(W)$, where the first equality follows from Theorem 6.3 and the second equality from Proposition 6.9. $\square$

Proposition 7.6. *Let $L(W^1)$, $L(W^2)$ be simple modules corresponding to cluster variables w_1 , w_2 (either via Proposition 7.5 or x_i , f_i). Then $L(W^1) \otimes L(W^2)$ is simple if and only if w_1 and w_2 live in a common cluster.*

For type ADE , this shows the condition (1) in the monoidal categorification 2.4.

Proof. The assertion is trivial when $W^1 = W^2$ by Corollary 6.16, since $\dim W^1 = \dim W^2$ is a real Schur root.

The assertion is also trivial for f_i by Proposition 6.7. So we may assume both W^1 and W^2 are not f_i . Therefore we have $W^1 = \varphi W^1$, $W^2 = \varphi W^2$.

By Propositions 6.10, 6.12 $L(W^1) \otimes L(W^2)$ is simple if and only if $\text{ext}^1(W^1, W^2) = \text{ext}^1(W^2, W^1) = 0$.

If both $L(W^1)$, $L(W^2)$ are not x_i nor x'_i , then $L(W^1) = z[\sigma W^1]$, $L(W^2) = z[\sigma W^2]$ as in the proof of Proposition 7.5. We have $\text{ext}^1(W^1, W^2) = \text{ext}^1(W^2, W^1) = 0$ if and only if $\text{ext}^1(\sigma W^1, \sigma W^2) = \text{ext}^1(\sigma W^2, \sigma W^1) = 0$. This happens if and only if $\sigma W^1 \oplus \sigma W^2$ is rigid, and hence can be extended to a tilting module. From §§7.2, 7.3, this is equivalent to that the corresponding cluster variables live in a common cluster.

If $L(W^1) = x_i$, $L(W^2) = x'_i$, then $L(W^1) \otimes L(W^2)$ is not simple by the T -system (5.2). They are not in any cluster simultaneously. Any other pairs from x_i , x'_j , they are always in a common cluster. It is also clear that $L(W^1) \otimes L(W^2)$ is always simple. Therefore we may assume $L(W^1)$ is one of x_i , x'_i , and $L(W^2)$ is not.

Consider the case $L(W^1) = x_i$ with $i \in I_0$. We have $W^1 = S_{i'}$. From Proposition 6.7, 6.8 $L(S_{i'}) \otimes L(W^2)$ is simple if and only if $W_i^2 = 0$. In this case $x_i = z_i$ is a cluster variable from the seed for $\mathbf{z}$ -quiver. From §7.2 the cluster variable w^2 is in a common cluster with z_i if and only if $\sigma W_i^2 = 0$. This is equivalent to $W_i^2 = 0$, since $i \in I_0$.

The case $L(W^1) = x'_i$ with $i \in I_0$ is not necessary to consider since we have $L(W^1) = L(S_i) = z[S_i]$ and already studied.

Next suppose $L(W^1) = x_i$ with $i \in I_1$. We have $W^1 = S_{i'}$. From Proposition 6.7, 6.8 $L(S_{i'}) \otimes L(W^2)$ is simple if and only if $W_i^2 = 0$ as before. Since i is a source, this is equivalent to $\text{Hom}(W^2, S_i) = 0$. From the definition of the reflection functor, it is equivalent to $\text{Ext}^1(S_i, \sigma W^2) = 0$. Since we have $x_i = z_i^*$, the corresponding rigid module for the $\mathbf{z}$ -quiver is S_i . Therefore x_i and w is in a common cluster if and only if $\text{Ext}^1(S_i, \sigma W^2) = 0 = \text{Ext}^1(\sigma W^2, S_i)$ by §7.2. But the latter equality is trivial since i is source. Thus we have checked the assertion in this case.

Finally suppose $L(W^1) = x'_i$ for $i \in I_1$. This is z_i and corresponds to a vertex i in the cluster-tilting set for $\mathbf{z}$ -quiver. Therefore w is in a same cluster with z_i if and only if $\sigma W_i^2 = 0$. By the same argument as above, this is equivalent to $\text{Ext}^1(S_i, W^2) = 0 = \text{Ext}^1(W^2, S_i)$. Thus we have checked the final case. $\square$

Remark 7.7. As indicated in the proof, it is more natural to define σS_i as $S_i[-1]$, an object in the derived category $\mathcal{D}(\text{rep } \sigma \tilde{\mathcal{Q}}^{\text{op}})$. This is also compatible with the cluster category theory, as $S_i[-1] = I_i[-1]$ for $i \in I_1$, where I_i is the indecomposable injective module corresponding to the vertex i .

7.6. Exchange relation. Consider an exchange relation (2.3). Thanks to Propositions 7.5, 7.6 we have the corresponding equality in $\mathbf{R}_{\ell=1}$:

$$L(x_k) \otimes L(x_k^*) = L(m_+) + L(m_-).$$

Since $L(m_{\pm})$ are simple, this inequality in the Grothendieck group implies either of the followings:

$$0 \rightarrow L(m_+) \rightarrow L(x_k) \otimes L(x_k^*) \rightarrow L(m_-) \rightarrow 0,$$

or

$$0 \rightarrow L(m_-) \rightarrow L(x_k) \otimes L(x_k^*) \rightarrow L(m_+) \rightarrow 0$$

in the level of modules. It is natural to conjecture that we always have the above one. For the T -system, this is true thanks to Remark 5.3.

This conjecture follows from a refinement of the exchange relation:

$$\chi_{q,t}(L(x_k) \otimes L(x_k^*)) = t^{-l+n} \chi_{q,t}(L(m_+)) + t^n \chi_{q,t}(L(m_-))$$

for some $l > 0, n \in \mathbb{Z}$. If we write the corresponding perverse sheaves by $P(x_k), P(x_k^*), P(m_+), P(m_-)$, the above means that

$$\begin{aligned} \text{Res}(P(m_+)) &= P(x_k) \boxtimes P(x_k^*)[l-n] \oplus \cdots, \\ \text{Res}(P(m_-)) &= P(x_k) \boxtimes P(x_k^*)[-n] \oplus \cdots, \end{aligned}$$

where $\cdots$ means sum of (shifts of) other perverse sheaves. Since $\text{Hom}(P(x_k) \boxtimes P(x_k^*)[l], P(x_k) \boxtimes P(x_k^*))$ vanishes for $l > 0$ by a property of perverse sheaves [12, 8.4.4], we see that $L(m_+)$ is a submodule of $L(x_k) \otimes L(x_k^*)$.

This refinement of the exchange relation might be proved directly, but it should be proved naturally if we make an isomorphism of the quantum cluster algebra [4] with $\mathbf{R}_{t,\ell=1}$.

REFERENCES

- [1] I. Assem, D. Simson, and A. Skowroński, *Elements of the representation theory of associative algebras. Vol. 1*, London Mathematical Society Student Texts, vol. 65, Cambridge University Press, Cambridge, 2006, Techniques of representation theory.
- [2] A. A. Beilinson, J. Bernstein, and P. Deligne, *Faisceaux pervers*, Analysis and topology on singular spaces, I (Luminy, 1981), Astérisque, vol. 100, Soc. Math. France, Paris, 1982, pp. 5–171.
- [3] A. Berenstein, S. Fomin, and A. Zelevinsky, *Cluster algebras. III. Upper bounds and double Bruhat cells*, Duke Math. J. **126** (2005), no. 1, 1–52.
- [4] A. Berenstein and A. Zelevinsky, *Quantum cluster algebras*, Adv. Math. **195** (2005), no. 2, 405–455.
- [5] I. N. Bernšteĭn, I. M. Gel’fand, and V. A. Ponomarev, *Coxeter functors, and Gabriel’s theorem*, Uspehi Mat. Nauk **28** (1973), no. 2(170), 19–33.
- [6] A. B. Buan, R. Marsh, M. Reineke, I. Reiten, and G. Todorov, *Tilting theory and cluster combinatorics*, Adv. Math. **204** (2006), no. 2, 572–618.
- [7] P. Caldero and F. Chapoton, *Cluster algebras as Hall algebras of quiver representations*, Comment. Math. Helv. **81** (2006), no. 3, 595–616.
- [8] P. Caldero and B. Keller, *From triangulated categories to cluster algebras. II*, Ann. Sci. École Norm. Sup. (4) **39** (2006), no. 6, 983–1009.
- [9] P. Caldero and M. Reineke, *On the quiver Grassmannian in the acyclic case*, J. Pure Appl. Algebra **212** (2008), no. 11, 2369–2380.
- [10] P. Caldero and A. Zelevinsky, *Laurent expansions in cluster algebras via quiver representations*, Mosc. Math. J. **6** (2006), no. 3, 411–429.
- [11] V. Chari and D. Hernandez, *Beyond Kirillov-Reshetikhin modules*, 2008, arXiv.org:0812.1716.
- [12] N. Chriss and V. Ginzburg, *Representation theory and complex geometry*, Birkhäuser Boston Inc., Boston, MA, 1997.
- [13] J. H. Conway and H. S. M. Coxeter, *Triangulated polygons and frieze patterns*, Math. Gaz. **57** (1973), no. 400, 87–94.
- [14] W. Crawley-Boevey, *Normality of Marsden-Weinstein reductions for representations of quivers*, Math. Ann. **325** (2003), no. 1, 55–79.
- [15] V. I. Danilov and A. G. Khovanskii, *Newton polyhedra and an algorithm for calculating Hodge-Deligne numbers*, Izv. Akad. Nauk SSSR Ser. Mat. **50** (1986), no. 5, 925–945.
- [16] P. Deligne, *Théorie de Hodge. III*, Inst. Hautes Études Sci. Publ. Math. (1974), no. 44, 5–77.
- [17] H. Derksen, J. Weyman, and A. Zelevinsky, *Quivers with potentials and their representations II: Applications to cluster algebras*, 2009, arXiv.org:0904.0676.
- [18] P. Di Francesco and R. Kedem, *Q-systems, heaps, paths and cluster positivity*, 2008, arXiv.org:0811.3027.
- [19] M. Ding, J. Xiao, and F. Xu, *Integral bases of cluster algebras and representations of tame quivers*, 2009, arXiv.org:0901.1937.

- [20] G. Dupont, *Generic variables in acyclic cluster algebras*, 2008, arXiv.org:0811.2909.
- [21] S. Fomin and A. Zelevinsky, *Cluster algebras. I. Foundations*, J. Amer. Math. Soc. **15** (2002), no. 2, 497–529 (electronic).
- [22] ———, *Cluster algebras. II. Finite type classification*, Invent. Math. **154** (2003), no. 1, 63–121.
- [23] ———, *Cluster algebras. IV. Coefficients*, Compos. Math. **143** (2007), no. 1, 112–164.
- [24] E. Frenkel and N. Reshetikhin, *The q -characters of representations of quantum affine algebras and deformations of $\mathcal{W}$ -algebras*, Recent developments in quantum affine algebras and related topics (Raleigh, NC, 1998), Contemp. Math., vol. 248, Amer. Math. Soc., Providence, RI, 1999, pp. 163–205.
- [25] C. Geiß, B. Leclerc, and J. Schröer, *Rigid modules over preprojective algebras*, Invent. Math. **165** (2006), no. 3, 589–632.
- [26] ———, *Cluster algebra structures and semicanonical bases for unipotent groups*, 2007, arXiv.org:math/0703039.
- [27] S. I. Gelfand and Y. I. Manin, *Methods of homological algebra*, second ed., Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2003.
- [28] D. Happel and L. Unger, *Almost complete tilting modules*, Proc. Amer. Math. Soc. **107** (1989), no. 3, 603–610.
- [29] D. Hernandez, *The Kirillov-Reshetikhin conjecture and solutions of T -systems*, J. Reine Angew. Math. **596** (2006), 63–87.
- [30] ———, *Smallness problem for quantum affine algebras and quiver varieties*, Ann. Sci. Éc. Norm. Supér. (4) **41** (2008), no. 2, 271–306.
- [31] D. Hernandez and B. Leclerc, *Cluster algebras and quantum affine algebras*, 2009, arXiv.org:0903.1452.
- [32] A. Hubery, *Acyclic cluster algebras via Ringel-Hall algebras*, preprint.
- [33] R. Inoue, O. Iyama, A. Kuniba, T. Nakanishi, and J. Suzuki, *Periodicities of T -systems and Y -systems*, 2008, arXiv.org:0812.06.
- [34] V. G. Kac, *Infinite root systems, representations of graphs and invariant theory*, Invent. Math. **56** (1980), no. 1, 57–92.
- [35] ———, *Infinite root systems, representations of graphs and invariant theory. II*, J. Algebra **78** (1982), no. 1, 141–162.
- [36] M. Kashiwara and P. Schapira, *Sheaves on manifolds*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 292, Springer-Verlag, Berlin, 1990, With a chapter in French by Christian Houzel.
- [37] B. Keller, *Cluster algebras, quiver representations and triangulated categories*, 2008, arXiv.org:0807.1960.
- [38] H. Knight, *Spectra of tensor products of finite-dimensional representations of Yangians*, J. Algebra **174** (1995), no. 1, 187–196.
- [39] G. Laumon, *Transformation de Fourier, constantes d'équations fonctionnelles et conjecture de Weil*, Inst. Hautes Études Sci. Publ. Math. (1987), no. 65, 131–210.
- [40] B. Leclerc, *Algèbres affine quantiques et algèbres amassées*, Jan. 2008, talk at IHP.
- [41] ———, *Canonical and semicanonical bases*, Nov. 2008, talk at Reims.
- [42] G. Lusztig, *Affine quivers and canonical bases*, Inst. Hautes Études Sci. Publ. Math. (1992), no. 76, 111–163.
- [43] ———, *Introduction to quantum groups*, Progress in Mathematics, vol. 110, Birkhäuser Boston Inc., Boston, MA, 1993.
- [44] ———, *On quiver varieties*, Adv. Math. **136** (1998), no. 1, 141–182.
- [45] ———, *Semicanonical bases arising from enveloping algebras*, Adv. Math. **151** (2000), no. 2, 129–139.
- [46] R. Marsh, M. Reineke, and A. Zelevinsky, *Generalized associahedra via quiver representations*, Trans. Amer. Math. Soc. **355** (2003), no. 10, 4171–4186 (electronic).
- [47] H. Nakajima, *Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras*, Duke Math. J. **76** (1994), no. 2, 365–416.
- [48] ———, *Quiver varieties and Kac-Moody algebras*, Duke Math. J. **91** (1998), no. 3, 515–560.
- [49] ———, *Quiver varieties and finite-dimensional representations of quantum affine algebras*, J. Amer. Math. Soc. **14** (2001), no. 1, 145–238 (electronic).
- [50] ———, *Quiver varieties and tensor products*, Invent. Math. **146** (2001), no. 2, 399–449.
- [51] ———, *T -analogue of the q -characters of finite dimensional representations of quantum affine algebras*, Physics and combinatorics, 2000 (Nagoya), World Sci. Publ., River Edge, NJ, 2001, pp. 196–219.

- [52] ———, *Geometric construction of representations of affine algebras*, Proceedings of the International Congress of Mathematicians, Vol. I (Beijing, 2002) (Beijing), Higher Ed. Press, 2002, pp. 423–438.
- [53] ———, *t-analogs of q-characters of Kirillov-Reshetikhin modules of quantum affine algebras*, Represent. Theory **7** (2003), 259–274 (electronic).
- [54] ———, *Quiver varieties and t-analogs of q-characters of quantum affine algebras*, Ann. of Math. (2) **160** (2004), no. 3, 1057–1097.
- [55] ———, *t-analogs of q-characters of quantum affine algebras of type E_6 , E_7 , E_8* , 2006, arXiv.org:math/0606637.
- [56] A. Schofield, *General representations of quivers*, Proc. London Math. Soc. (3) **65** (1992), no. 1, 46–64.
- [57] P. Sherman and A. Zelevinsky, *Positivity and canonical bases in rank 2 cluster algebras of finite and affine types*, Mosc. Math. J. **4** (2004), no. 4, 947–974, 982.
- [58] M. Varagnolo and E. Vasserot, *Perverse sheaves and quantum Grothendieck rings*, Studies in memory of Issai Schur (Chevaleret/Rehovot, 2000), Progr. Math., vol. 210, Birkhäuser Boston, Boston, MA, 2003, pp. 345–365.

RESEARCH INSTITUTE FOR MATHEMATICAL SCIENCES, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN

E-mail address: nakajima@math.kyoto-u.ac.jp

URL: <http://www.math.kyoto-u.ac.jp/~nakajima>